


UTMS 2006–9

May 8, 2006

**On the Young theorem
for amalgams and Besov spaces**

by

Yoshihiro SAWANO and Tsuyoshi YONEDA



UNIVERSITY OF TOKYO

GRADUATE SCHOOL OF MATHEMATICAL SCIENCES

KOMABA, TOKYO, JAPAN

On the Young theorem for amalgams and Besov spaces

May 8, 2006

*Yoshihiro Sawano and Tsuyoshi Yoneda**
Graduate School of Mathematical Sciences, The University of Tokyo,
3-8-1 Komaba, Meguro-ku Tokyo 153-8914, JAPAN
E-mail: yoshihiro@ms.u-tokyo.ac.jp yoneda@ms.u-tokyo.ac.jp

Abstract

In this paper, we obtain a refinement of the Young theorem. The Young theorem tells us that the Fourier transform \mathcal{F} sends the L^p functions to the $L^{p'}$ functions, if $1 \leq p \leq 2$. This theorem has a refinement. For example, $\mathcal{F} : L^1 \rightarrow B_{\infty 1}^0$, where B_{pq}^s is the Besov space. In this present paper we shall consider the more refined version of this theorem by using the amalgams and the Besov spaces.

Keywords Besov space, amalgam space, Fourier transform

2000 Mathematics Subject Classification Primary 42B10; Secondary 42B35.

1 Introduction

The aim of this paper is to refine the Young theorem. The Young theorem, as is well-known, asserts that the range of the L^p space by the Fourier transform is the $L^{p'}$ space, whenever $1 \leq p \leq 2$.

$$\mathcal{F} : L^p \rightarrow L^{p'}.$$

Here and below, for definiteness, we define the Fourier transform of $f \in L^1 \cap L^p$ to be

$$\mathcal{F}f(\xi) := \int_{\mathbf{R}^n} f(x) e^{-i\xi \cdot x} dx.$$

The above well-known theorem has a following refinement.

$$\mathcal{F} : L^p \rightarrow B_{p'p}^0,$$

* The authors are supported by Research Fellowships of the Japan Society for the Promotion of Science for Young Scientists.

where B_{pq}^s is the Besov space whose definition will be recalled later. It is known that $B_{\infty 1}^0$ is a function space contained in L^∞ . For some related facts we refer to [1, p164]. Let C denote the space of all bounded continuous functions. Since $B_{\infty 1}^0 \subset C$, the Besov space $B_{\infty 1}^0$ describes the situation more precisely than the Lebesgue space L^∞ . However, once the Besov spaces comes into the play, we hit upon a natural question. How can we make use of the information of q in the Besov spaces B_{pq}^s ?

In this present paper we give a more refined version of this theorem and show the sharpness of our result. In Section 2 we give the definition of function spaces to formulate our theorem. The proof of the theorem is contained in Section 3. Finally in Section 4 we exhibit examples showing the sharpness of our result.

2 Function spaces

In this section we present the definition of function spaces we work on.

Besov space Following [4, 5], we give a definition of Besov spaces. The definition is somehow different from those in [4, 5]. However the resulting norms will be equivalent. We use \mathbf{N}_0 to denote $\{0, 1, 2, \dots\}$.

First, given a complex sequence $\{a_j\}_{j \in \mathbf{N}_0}$, we set $\|a_j : l^q\| := \left(\sum_{j \in \mathbf{N}_0} |a_j|^q \right)^{\frac{1}{q}}$, $0 < q \leq \infty$.

We also define $\|a_z : l^q\| := \left(\sum_{z \in \mathbf{Z}^n} |a_z|^q \right)^{\frac{1}{q}}$, $0 < q \leq \infty$ for $\{a_z\}_{z \in \mathbf{Z}^n}$. If possible confusion can occur, we write

$$\|\{a_j\}_j : l^q\|, \|\{a_z\}_z : l^q\|$$

instead of $\|a_j : l^q\|$, $\|a_z : l^q\|$. Similarly, the notation $\|\{a_{j,z}\}_{j,z} : l^q\|$ means the l^q -norm of $\{a_{j,z}\}_{j \in \mathbf{N}_0, z \in \mathbf{Z}^n}$. Next, for a sequence of complex valued measurable functions $\{f_j\}_{j \in \mathbf{N}_0}$, we set

$$\|f_j : l^q(L^p)\| := \|\|f_j : L^p\| : l^q\|, \quad 0 < p, q \leq \infty.$$

If $p = \infty$ and / or $q = \infty$, we make a natural modification in the above formulae.

Definition 2.1. Let $\phi_0, \phi_1 \in \mathcal{S}$ be even functions satisfying the following conditions.

$$\chi_{[-2,2]^n} \leq \phi_0 \leq \chi_{[-4,4]^n}, \quad \chi_{[-4,4]^n \setminus [-2,2]^n} \leq \phi_1 \leq \chi_{[-8,8]^n \setminus [-1,1]^n}.$$

We set $\phi_j(x) := \phi_1(2^{-j+1}x)$ for $j \geq 2$. For $f \in \mathcal{S}'$, we denote $\phi_j(D)f := \mathcal{F}^{-1}(\phi_j \cdot \mathcal{F}f)$.

What counts about this definition is to adopt cubes instead of balls. We prefer to use cubes because we will consider the amalgam spaces. With this preparation in mind, we shall define the Besov norms.

Definition 2.2. Let $0 < p, q \leq \infty$ and $s \in \mathbf{R}$. Under the notations in Definition 2.1, we define B_{pq}^s to be the set of the Schwartz distributions $f \in \mathcal{S}'$ for which the quasi-norm

$$\|f : B_{pq}^s\| := \|2^{js} \phi_j(D)f : l^q(L^p)\| \quad (1)$$

is finite.

It can be easily shown that the definition of the Besov space B_{pq}^s is independent of the choice of ϕ_0, ϕ_1 by virtue of [4, Theorem 1.6.3].

(Weighted) amalgam space Now we will follow [2, 3] for the definitions. Given a measurable set A , we set

$$\|f : L^p(A)\| := \|\chi_A f : L^p\|.$$

Definition 2.3. Let $0 < p, q \leq \infty$ and $s \in \mathbf{R}$. Set $Q_z := z + [0, 1]^n$ for $z \in \mathbf{Z}^n$, the translation of the unit cube. For a Lebesgue locally integrable function f we define

$$\|f : (L^p, l^q(\langle z \rangle^s))\| := \|\langle z \rangle^s \cdot \|f : L^p(Q_z)\| : l^q\|,$$

where $\langle a \rangle := \sqrt{|a|^2 + 1}$ for $a \in \mathbf{R}^n$. $(L^p, l^q(\langle z \rangle^s))$ is a set of all locally integrable functions f for which the quasi-norm $\|f : (L^p, l^q(\langle z \rangle^s))\| < \infty$. For brevity we write $(L^p, l^q) := (L^p, l^q(1))$.

It can be seen that $(L^p, l^p) = L^p$ with norm coincidence. By definition of the norm, the following multiplication operator is an isomorphism.

$$f \in (L^p, l^q(\langle z \rangle^s)) \mapsto \langle \cdot \rangle^t \cdot f \in (L^p, l^q(\langle z \rangle^{s-t})). \quad (2)$$

Note that $(L^p, l^q(\langle z \rangle^s)) \subset \mathcal{S}'$, if $1 \leq p \leq \infty$, $0 < q \leq \infty$ and $s \in \mathbf{R}$. It can be easily seen that

$$(L^{p_1}, l^{q_1}(\langle z \rangle^{s_1})) \subset (L^{p_2}, l^{q_2}(\langle z \rangle^{s_2}))$$

for $p_1 \geq p_2$, $q_1 \leq q_2$ and $s_1 \geq s_2$.

Main theorem With these definitions in mind, we formulate our main theorem.

Theorem 2.4. 1. Let $1 \leq p \leq 2$, $0 < q \leq \infty$ and $s \in \mathbf{R}$. Then

$$\mathcal{F} : (L^p, l^q(\langle z \rangle^s)) \rightarrow B_{p'q}^{s-n(\frac{1}{p}-\frac{1}{q})_+}. \quad (3)$$

2. Let $1 \leq p \leq 2$, $0 < q \leq \infty$ and $s \in \mathbf{R}$. Then

$$\mathcal{F} : B_{pq}^s \rightarrow \left(L^{p'}, l^q(\langle z \rangle^{s-n(\frac{1}{q}-\frac{1}{p'})_+}) \right). \quad (4)$$

Here and below, for $a \in \mathbf{R}$ we write $a_+ := \max(a, 0)$.

Before we come to the proof, we state one more corollary.

Corollary 2.5. Suppose that $2 \leq p \leq \infty$, $0 < q \leq \infty$ and $s \in \mathbf{R}$. Then

$$\mathcal{F} : (L^p, l^q(\langle z \rangle^s)) \rightarrow B_{2q}^{s-n(\frac{1}{2}-\frac{1}{q})_+}.$$

In particular,

$$\mathcal{F} : L^\infty \rightarrow B_{2\infty}^{-\frac{n}{2}}.$$

Once we obtain Theorem 2.4, Corollary 2.5 is easy to prove: All we have to note is

$$(L^p, l^q(\langle z \rangle^s)) \subset (L^2, l^q(\langle z \rangle^s))$$

for $2 \leq p \leq \infty$.

The rest of this paper is devoted to the proof of Theorem 2.4 and to investigating the sharpness of these results.

3 Proof of Theorem 2.4

First, we recall the boundedness of the lift operator.

$$(1 - \Delta)^{\frac{s}{2}} : B_{pq}^s \rightarrow B_{pq}^{s-t}. \quad (5)$$

For the proof of (3), (2) and (5) allow us to assume $s = 0$.

With this preparation our present task is to estimate

$$\left\| 2^{-jn(\frac{1}{p} - \frac{1}{q})_+} \phi_j(D) \mathcal{F}f : l^q(L^{p'}) \right\| \quad (6)$$

for $f \in (L^p, l^q(\langle z \rangle^s))$. Note that the Young theorem gives

$$\|\phi_j(D) \mathcal{F}f : L^{p'}\| \leq \|\phi_j \cdot \mathcal{F}(\mathcal{F}f) : L^p\|.$$

Set $A_j := \text{supp}(\phi_j)$ for $j \in \mathbf{N}_0$. Then (6) can be majorized by

$$\left\| 2^{-jn(\frac{1}{p} - \frac{1}{q})_+} \|\mathcal{F}(\mathcal{F}f) : L^p(A_j)\| : l^q \right\|.$$

Since $\mathcal{F}(\mathcal{F}f)(x) = cf(-x)$ and the ϕ_j are even, we have

$$\|\mathcal{F}(\mathcal{F}f) : L^p(A_j)\| = c \|f : L^p(A_j)\|. \quad (7)$$

The inequality $\left(\sum_{j=1}^N |a_j| \right)^{\frac{q}{p}} \leq N^{\left(\frac{q}{p}-1\right)_+} \sum_{j=1}^N |a_j|^{\frac{q}{p}}$ gives us

$$\|f : L^p(A_j)\|^q \leq c 2^{jqn(\frac{1}{p} - \frac{1}{q})_+} \sum_{z \in \mathbf{Z}^n} \|f : L^p(A_j \cap Q_z)\|^q. \quad (8)$$

If we put (7) and (8) together, then we have

$$\left\| 2^{-jn(\frac{1}{p} - \frac{1}{q})_+} \phi_j(D) \mathcal{F}f : l^q(L^{p'}) \right\| \leq c \left\| \left\{ \|f : L^p(A_j \cap Q_z)\| \right\}_{j,z} : l^q \right\|. \quad (9)$$

Given $z \in \mathbf{Z}^n$, from the definition of the A_j , there are at most three j such that $A_j \cap Q_z \neq \emptyset$, and hence,

$$\sum_{j \in \mathbf{N}_0} \|f : L^p(A_j \cap Q_z)\|^q \leq 3 \|f : L^p(Q_z)\|^q. \quad (10)$$

Combining (9) and (10), we obtain

$$\left\| 2^{-jn(\frac{1}{p} - \frac{1}{q})_+} \phi_j(D) \mathcal{F}f : l^q(L^{p'}) \right\| \leq c \left\| \|f : L^p(Q_z)\| : l^q \right\| = \|f : (L^p, l^q)\|.$$

This is the desired result.

Next, we prove (4). As before, we assume $s = 0$. Let $|z|_\infty$ denote $\max(|z_1|, |z_2|, \dots, |z_n|)$. First, we observe by the definition of the norm,

$$\left\| \mathcal{F}f : \left(L^{p'}, l^q \left(\langle z \rangle^{-n(\frac{1}{q} - \frac{1}{p'})_+} \right) \right) \right\| \sim \left\| \left\{ 2^{-jn(\frac{1}{q} - \frac{1}{p'})_+} \sum_{\substack{z \in \mathbf{Z}^n \\ [2^{j-1}] \leq |z|_\infty < 2^j}} \|\mathcal{F}f : L^{p'}(Q_z)\| \right\}_j : l^q \right\|,$$

where $[\cdot]$ denotes the Gauss sign.

The inequality $\sum_{j=1}^N |a_j|^{\frac{q}{p'}} \leq N^{(1-\frac{q}{p'})_+} \left(\sum_{j=1}^N |a_j| \right)^{\frac{q}{p'}}$ and Young's theorem yield

$$\begin{aligned} \left\| \mathcal{F}f : \left(L^{p'}, l^q \left(\langle z \rangle^{-n \left(\frac{1}{q} - \frac{1}{p'} \right)_+} \right) \right) \right\| &\leq c \left\| \left\{ \left(\sum_{\substack{z \in \mathbf{Z}^n \\ [2^{j-1}] \leq |z|_\infty < 2^j}} \|\mathcal{F}f : L^{p'}(Q_z)\|^{p'} \right)^{\frac{1}{p'}} \right\}_j : l^q \right\| \\ &\sim \left\| \phi_j \cdot \mathcal{F}f : l^q(L^{p'}) \right\| = c \left\| \mathcal{F}\phi_j(D)f : l^q(L^{p'}) \right\| \leq c \left\| \phi_j(D)f : l^q(L^p) \right\| = c \|f : B_{pq}^0\|. \end{aligned}$$

This proves (4).

4 Sharpness of Theorem 2.4

In this section we deduce some necessary conditions. We consider the following problem :

Problem 4.1. *Let $1 \leq p_1 \leq \infty$, $0 < p_2 \leq \infty$, $0 < q_1, q_2 \leq \infty$ and $s_1, s_2 \in \mathbf{R}$. Under what condition does the Fourier transform \mathcal{F} send $(L^{p_1}, l^{q_1}(\langle z \rangle^{s_1}))$ continuously to $B_{p_2 q_2}^{s_2}$? That is, when is the estimate*

$$\|\mathcal{F}f : B_{p_2 q_2}^{s_2}\| \leq c \|f : (L^{p_1}, l^{q_1}(\langle z \rangle^{s_1}))\| \quad (11)$$

is true ? Find necessary conditions of (11).

First, we prove that the smoothness parameter s cannot be improved.

Proposition 4.2. *If (11) is true, then $s_2 \leq s_1$.*

Proof. Let $\tau \in \mathcal{S}$ be an even function with $\chi_{B(1/4)} \leq \tau \leq \chi_{B(1/2)}$, where $B(r)$ denotes the open ball centered at the origin of radius $r > 0$. We set $e_1 = (1, 0, 0, \dots, 0)$, the elementary vector in \mathbf{R}^n , and define $\tau_j(x) := \tau(x - 2^j e_1)$. Then we obtain

$$\|\tau_j : (L^{p_1}, l^{q_1}(\langle z \rangle^{s_1}))\| \sim 2^{js_1}, \quad \|\mathcal{F}\tau_j : B_{p_2 q_2}^{s_2}\| \sim \|2^{js_2} \mathcal{F}\tau_j : L^{p_2}\| \sim \|2^{js_2} \mathcal{F}\tau : L^{p_2}\| \sim 2^{js_2}.$$

Since by assumption we have $\|\mathcal{F}f : B_{p_2 q_2}^{s_2}\| \leq c \|f : (L^{p_1}, l^{q_1}(\langle z \rangle^{s_1}))\|$ for all $f \in (L^{p_1}, l^{q_1}(\langle z \rangle^{s_1}))$, it follows that $s_2 \leq s_1$. ■

Next, we discuss how the integrability parameter changes by the Fourier transform.

Proposition 4.3. *If (11) is true, then $p_2 \geq p_1'$.*

Proof. We use τ in the proof in Proposition 4.2 again. Consider $f(x) := |x|^{-\alpha} \tau(x)$. Set $\delta = 1 - \tau$ and $g(x) := |x|^{-\alpha} \delta(x)$. It is well-known that

$$\mathcal{F}f(\xi) + \mathcal{F}g(\xi) = c |\xi|^{\alpha-n}.$$

Since $|\xi|^{2N} \mathcal{F}g(\xi) = \mathcal{F}[(-\Delta)^N g](\xi)$ and $(-\Delta)^N g \in L^1$ for $N \gg 1$, it follows that $|\mathcal{F}g(\xi)| \leq c |\xi|^{-2N}$. From this we deduce

$$|\mathcal{F}f(\xi)| \sim |\xi|^{\alpha-n} \text{ as } \xi \rightarrow \infty.$$

$f \in (L^{p_1}, l^{q_1}(\langle z \rangle^{s_1}))$ if and only if $p_1 \alpha < n$. Meanwhile, from the definition of the norm, $\mathcal{F}f \in B_{p_2 q_2}^{s_2}$ forces $p_2(n - \alpha) > n$. Thus it is necessary that $\alpha < n/p_1$ implies $\alpha < n - n/p_2$. From this it follows that $n/p_1 \leq n - n/p_2$, which is equivalent to $p_2 \geq p_1'$. ■

The restrictions appearing in Propositions 4.2 and 4.3 are natural, if we take into account $\mathcal{F} : L^p(\langle z \rangle^s) \rightarrow W_{p'}^s$, where $W_{p'}^s$ denotes the Sobolev space. It is well-known that

$$B_{pq}^{s+\varepsilon} \subset B_{p'q'}^s, \quad 0 < p, q, q' \leq \infty, \quad \varepsilon > 0$$

and

$$B_{p_1 q}^{s_1} \subset B_{p_2 q}^{s_2}, \quad 0 < p_1, p_2, q \leq \infty, \quad s_1 > s_2, \quad s_1 - \frac{n}{p_1} = s_2 - \frac{n}{p_2}. \quad (12)$$

Thus, if $s_1 > s_2$ or $p_1' < p_2$, then the situation can be considered degenerate. Next, we explain the decay of the parameter s in Theorem 2.4 when $p < q$.

Proposition 4.4. *If (11) is true, then $s_2 \leq s_1 - \frac{n}{p_2'} + \frac{n}{q_1}$.*

From this proposition, the restriction $s_2 \leq s_1 - \frac{n}{p_2'} + \frac{n}{q_1}$ in Theorem 2.4 is essential.

Proof. Let ϕ_j be the function in Definition 2.2.

$$\|\phi_j : (L^{p_1}, l^{q_1}(\langle z \rangle^{s_1}))\| \sim 2^j \left(s_1 + \frac{n}{q_1} \right) \quad \text{and} \quad \|\mathcal{F}\phi_j : B_{p_2 q_2}^{s_2}\| \sim \|2^{j s_2} \mathcal{F}\phi_j : L^{p_2}\| \sim 2^j \left(s_2 + \frac{n}{p_2'} \right).$$

As a result the desired inequality follows. ■

We tackle a subtler problem: Can we improve q_2 in (11)? The case when $s_2 < \min(s_1, s_1 - n/p_2' + n/q_1)$ can be regarded as degenerate and we concentrate on the limit case.

Proposition 4.5. *Assume $s_2 = \min\left(s_1, s_1 - \frac{n}{p_2'} + \frac{n}{q_1}\right)$. If (11) is true, then $q_2 \geq q_1$.*

Proof. We consider two cases separately: $s_2 = s_1 - \frac{n}{p_2'} + \frac{n}{q_1}$ and $s_2 = s_1$. By using the lift operators, we may assume $s_1 = 0$.

First, we tackle the case $s_2 = s_1 - \frac{n}{p_2'} + \frac{n}{q_1}$. Let $\{a_j\}_{j \in \mathbf{N}}$ be a complex sequence as before such that $a_j = 0$ if 3 does not divide j . Let $f := \sum_{j \in \mathbf{N}_0} a_j \phi_j$, where the ϕ_j are from Definition 2.2. Then by the same reasoning as before we obtain

$$\|f : (L^{p_1}, l^{q_1})\| \sim \|2^{\frac{jn}{q_1}} a_j : l^{q_1}\| \quad \text{and} \quad \|\mathcal{F}f : B_{p_2 q_2}^{s_2}\| \sim \|2^{\frac{jn}{q_1}} a_j : l^{q_2}\|.$$

From this we deduce $q_2 \geq q_1$.

Now we turn to the case when $s_2 = s_1 = 0$. Then we use the τ_j in Proposition 4.2. Let $f := \sum_{j \in \mathbf{N}_0} a_j \tau_j$, where $\{a_j\}_{j \in \mathbf{N}_0}$ is a complex sequence as before. Then we have

$$\|f : (L^{p_1}, l^{q_1})\| \sim \|a_j : l^{q_1}\| \quad \text{and} \quad \|\mathcal{F}f : B_{p_2 q_2}^0\| \sim \|a_j : l^{q_2}\|.$$

Thus, we obtain $q_2 \geq q_1$. ■

Finally we show

Proposition 4.6. *Let $0 < q < 2$. Then the mapping $\mathcal{F} : (L^2, l^q) \rightarrow B_{2q}^0$ is not a surjection.*

Proof. By interpolation, we may assume that $1 \leq q < 2$. Assume that $\mathcal{F} : (L^2, l^q) \rightarrow B_{2q}^0$ is surjective. Then by duality we would have $\mathcal{F} : B_{2q'}^0 \rightarrow (L^2, l^{q'})$ is bijective. This would imply $\mathcal{F} : (L^2, l^{q'}) \rightarrow B_{2q'}^0$ is also bijective. This contradicts to Proposition 4.4. ■

Acknowledgement

The authors are very grateful to Prof. E. Nakai for his helpful discussion with them.

References

- [1] J. Bergh, L. Jörge, *Interpolation spaces. An introduction. Grundlehren der Mathematischen Wissenschaften, no. 223.* Springer-Verlag, Berlin-New York, 1976. Interpolation.
- [2] Finbarr Holland, *Harmonic analysis on amalgams*, J. London Math. Soc. (2), 10 (1975), 295-305.
- [3] John J. F. Fournier, Vancouver, *On the Hausdorff-Young Theorem for amalgams*, Mh. Math. 95, 117-135 (1983).
- [4] H. Triebel, *Theory of function spaces*, Birkhauser (1983).
- [5] H. Triebel, *Theory of function spaces II*, Birkhauser (1992).

UTMS

- 2005–46 Li Shumin and Masahiro Yamamoto: *An inverse problem for Maxwell's equations in anisotropic media.*
- 2005–47 Li Shumin: *Estimation of coefficients in a hyperbolic equation with impulsive inputs.*
- 2005–48 Mourad Bellassoued and Masahiro Yamamoto: *Lipschitz stability in determining density and two Lamé coefficients.*
- 2006–1 Takao Satoh: *Twisted second homology groups of the automorphism group of a free group.*
- 2006–2 Matthias M. Eller and Masahiro Yamamoto: *A Carleman inequality for the stationary anisotropic Maxwell system.*
- 2006–3 Mourad Choulli, Oleg Yu. Imanuvilov and Masahiro Yamamoto: *Inverse source problem for the Navier-Stokes equations.*
- 2006–4 Takashi Taniguchi: *Distributions of discriminants of cubic algebras.*
- 2006–5 Jin Cheng, Yanbo Wang and Masahiro Yamamoto: *Regularity of the Tikhonov regularized solutions and the exact solution.*
- 2006–6 Keiichi Gunji: *The dimension of the space of siegel Eisenstein series of weight one.*
- 2006–7 Takehiro Fujiwara: *Sixth order methods of Kusuoka approximation.*
- 2006–8 Takeshi Ohtsuka: *Motion of interfaces by the Allen-Cahn type equation with multiple-well potentials.*
- 2006–9 Yoshihiro Sawano and Tsuyoshi Yoneda: *On the Young theorem for amalgams and Besov spaces.*

The Graduate School of Mathematical Sciences was established in the University of Tokyo in April, 1992. Formerly there were two departments of mathematics in the University of Tokyo: one in the Faculty of Science and the other in the College of Arts and Sciences. All faculty members of these two departments have moved to the new graduate school, as well as several members of the Department of Pure and Applied Sciences in the College of Arts and Sciences. In January, 1993, the preprint series of the former two departments of mathematics were unified as the Preprint Series of the Graduate School of Mathematical Sciences, The University of Tokyo. For the information about the preprint series, please write to the preprint series office.

ADDRESS:

Graduate School of Mathematical Sciences, The University of Tokyo
3–8–1 Komaba Meguro-ku, Tokyo 153-8914, JAPAN
TEL +81-3-5465-7001 FAX +81-3-5465-7012