


UTMS 2010–16

October 25, 2010

**Uniform estimate for distributions
of the sum of i.i.d random variables
with fat tail: threshold case**

by

Kenji NAKAHARA



UNIVERSITY OF TOKYO

GRADUATE SCHOOL OF MATHEMATICAL SCIENCES

KOMABA, TOKYO, JAPAN

Uniform Estimate for Distributions of the Sum of i.i.d Random Variables with Fat Tail: Threshold case

Kenji NAKAHARA

Abstract

We show uniform estimates of distributions of the sum of i.i.d random variables in the threshold case. Rozovskii showed several uniform estimates but the speeds of convergence were not shown. Our main uniform estimate has a speed of convergence. We also compare our estimates with Nagaev's estimate which is valid in the non-threshold case and we show a necessary and sufficient condition for holding Nagaev's estimate in the threshold case.

1 Introduction

Let (Ω, \mathcal{F}, P) be a probability space and $X_n, n = 1, 2, \dots$, be independent identically distributed random variables whose probability law is μ . Let $F : \mathbb{R} \rightarrow [0, 1]$ and $\bar{F} : \mathbb{R} \rightarrow [0, 1]$ be given by $F(x) = \mu((-\infty, x])$ and $\bar{F}(x) = \mu((x, \infty))$, $x \in \mathbb{R}$. We assume the following.

(A1) $\bar{F}(x)$ is a regularly varying function of index $-\alpha$ for some $\alpha \geq 2$, as $x \rightarrow \infty$, i.e., if we let

$$L(x) = x^\alpha \bar{F}(x), \quad x \geq 1,$$

then $L(x) > 0$ for any $x \geq 1$, and for any $a > 0$

$$\frac{L(ax)}{L(x)} \rightarrow 1, \quad x \rightarrow \infty.$$

(A2) $\int_{-\infty}^0 |x|^{\alpha+\delta_0} \mu(dx) < \infty$ for some $\delta_0 \in (0, 1)$, $\int_{\mathbb{R}} x^2 \mu(dx) = 1$ and $\int_{\mathbb{R}} x \mu(dx) = 0$

Nagaev [3] proved the following theorem.

Theorem 1. (Nagaev) *Assume (A1) for $\alpha > 2$ and (A2). Then we have*

$$\sup_{s \in [1, \infty)} \left| \frac{P(\sum_{k=1}^n X_k > n^{1/2}s)}{\Phi_0(s) + n\bar{F}(n^{1/2}s)} - 1 \right| \rightarrow 0, \quad n \rightarrow \infty. \quad (1)$$

Here $\Phi_0 : \mathbb{R} \rightarrow \mathbb{R}$ is given by

$$\Phi_0(x) = \frac{1}{\sqrt{2\pi}} \int_x^\infty \exp\left(-\frac{y^2}{2}\right) dy, \quad x \in \mathbb{R}.$$

In this paper, we assume (A1) for $\alpha = 2$ (threshold case), (A2) and the following.

(A3) The probability law μ is absolutely continuous and has a density function $\rho : \mathbb{R} \rightarrow [0, \infty)$ which is right continuous and has a finite total variation.

Let us define $\Phi_k : \mathbb{R} \rightarrow \mathbb{R}$, $k = 1, 2, 3$, by

$$\Phi_1(x) = \frac{1}{\sqrt{2\pi}} \exp\left(-\frac{x^2}{2}\right) = -\frac{d}{dx} \Phi_0(x),$$

and

$$\Phi_k(x) = (-1)^{k-1} \frac{d^{k-1}}{dx^{k-1}} \Phi_1(x), \quad k = 2, 3.$$

Let $v_n = \int_{-\infty}^{n^{1/2}} x^2 \mu(dx)$ for $n \geq 1$ and let

$$\begin{aligned} & H(n, s) \\ &= \Phi_0(s) + n \int_{-\infty}^s \bar{F}((s-x)v_n^{1/2}n^{1/2}) \Phi_1(x) dx \\ & \quad - \left(v_n^{-1/2} n^{1/2} \Phi_1(s) \int_0^\infty x \mu(dx) + v_n^{-1} \frac{\Phi_2(s)}{2} \int_0^{n^{1/2}} x^2 \mu(dx) \right) \end{aligned}$$

Our main result is the following.

Theorem 2. *Assume (A1) for $\alpha = 2$, (A2) and (A3). Then for any $\delta \in (0, 1)$, there is a $C > 0$ such that*

$$\sup_{s \in [1, \infty)} \left| \frac{P(\sum_{k=1}^n X_k > n^{1/2}s)}{H(n, v_n^{-1/2}s)} - 1 \right| \leq CL(n^{1/2})^{1-\delta}, \quad n \geq 1. \quad (2)$$

In particular, we have

$$\sup_{s \in [1, \infty)} \left| \frac{P(\sum_{k=1}^n X_k > n^{1/2}s)}{\Phi_0(v_n^{-1/2}s) + n\bar{F}(n^{1/2}s)} - 1 \right| \rightarrow 0, \quad n \rightarrow \infty. \quad (3)$$

Rozovskii [4] showed different types of uniform estimate. (see Theorem 1, 2 and 3b in [4].) The estimates in Theorem 1, 2 in [4] were proved under more general assumptions but the speeds of convergence were not shown. The estimate in Theorem 3b in [4] is strongly related to Equation (3) but does not completely include our result. The proof of uniform estimates in [4] is different from ours.

We also prove the following.

Theorem 3. Assume (A1) for $\alpha = 2$, (A2) and (A3).

If $\limsup_{n \rightarrow \infty} (1 - v_n) \log \frac{1}{L(n^{1/2})} = 0$, then we have

$$\sup_{s \in [1, \infty)} \left| \frac{\Phi_0(v_n^{-1/2}s) + n\bar{F}(n^{1/2}s)}{\Phi_0(s) + n\bar{F}(n^{1/2}s)} - 1 \right| \rightarrow 0, \quad n \rightarrow \infty. \quad (4)$$

If $\limsup_{n \rightarrow \infty} (1 - v_n) \log \frac{1}{L(n^{1/2})} > 0$, then Equation (4) does not hold.

Combining with Theorem 2, Theorem 3 gives a necessary and sufficient condition for holding Equation (1). The condition $\limsup_{n \rightarrow \infty} (1 - v_n) \log \frac{1}{L(n^{1/2})} = 0$ is correspond to Equation (56) in [4]. Hence the estimate with $B_n = n^{1/2}$ in Theorem 3b in [4] is not valid under our assumptions.

We also prove the following to obtain Theorem 2.

Theorem 4. Assume (A1) for $\alpha = 2$, (A2) and (A3). Then for any $\delta \in (0, 1)$, there is a $C > 0$ such that

$$\left| P\left(\sum_{k=1}^n X_k > sn^{1/2}\right) - H(n, v_n^{-1/2}s) \right| \leq CL(n^{1/2})^{2-\delta}, \quad s \geq 1.$$

Thoughtout this paper we assume (A1) for $\alpha = 2$, (A2) and (A3). Then we see that $L(t) \rightarrow 0$, $t \rightarrow \infty$ and $\frac{1 - v_n}{L(n^{1/2})} \rightarrow \infty$, $n \rightarrow \infty$ (see Equation (5), (6)).

2 Preliminary facts

We summerize several known facts (c.f. Fushiya-Kusuoka[2]).

Proposition 1. We have

$$\sup_{1/2 \leq a \leq 2} \frac{L(ax)}{L(x)} \rightarrow 1, \quad x \rightarrow \infty,$$

and

$$\inf_{1/2 \leq a \leq 2} \frac{L(ax)}{L(x)} \rightarrow 1, \quad x \rightarrow \infty.$$

Proposition 2. For any $\varepsilon \in (0, 1)$, there is an $M(\varepsilon) \geq 1$ such that

$$M(\varepsilon)^{-1}y^{-\varepsilon} \leq \frac{L(yx)}{L(x)} \leq M(\varepsilon)y^\varepsilon \quad x, y \geq 1.$$

Proposition 3. (1) For any $\beta < -1$,

$$\frac{1}{t^{\beta+1}L(t)} \int_t^\infty x^\beta L(x) dx \rightarrow -\frac{1}{\beta+1}, \quad t \rightarrow \infty.$$

(2) For any $\beta > -1$,

$$\frac{1}{t^{\beta+1}L(t)} \int_1^t x^\beta L(x) dx \rightarrow \frac{1}{\beta+1}, \quad t \rightarrow \infty.$$

(3) Let $f : [1, \infty) \rightarrow (0, \infty)$ be given by

$$f(t) = \int_1^t x^{-1} L(x) dx \quad t \geq 1.$$

Then f is slowly varying. Moreover if $\lim_{t \rightarrow \infty} f(t) < \infty$, we have

$$\frac{1}{L(t)} \int_t^\infty x^{-1} L(x) dx \rightarrow \infty, \quad t \rightarrow \infty.$$

Proposition 4. There is a $C_0 > 0$ such that

$$|\Phi_k(x)| \leq C_0(1+x)^{k-1} \Phi_1(x), \quad x \geq 0, k = 1, 2,$$

and

$$C_0^{-1} \Phi_1(x) \leq x \Phi_0(x) \leq C_0 \Phi_1(x), \quad x \geq 1/2.$$

Proposition 5. (1) For any $m \geq 1$, let $r_{e,m} : \mathbb{R} \rightarrow \mathbb{C}$ be given by

$$r_{e,m}(t) = \exp(it) - \left(1 + \sum_{k=1}^m \frac{(it)^k}{k!}\right), \quad t \in \mathbb{R}.$$

Then we have

$$|r_{e,m}(t)| \leq \frac{\min(|t|^{m+1}, 2(m+1)|t|^m)}{(m+1)!}, \quad t \in \mathbb{R}.$$

(2) For any $m \geq 1$, let $r_{l,m} : \{z \in \mathbb{C}; |z| \leq 1/2\} \rightarrow \mathbb{C}$ be given by

$$r_{l,m}(z) = \log(1+z) - \sum_{k=1}^m \frac{(-1)^{k-1}}{k} z^k, \quad z \in \mathbb{C}, |z| \leq 1/2.$$

Then we have

$$|r_{l,m}(z)| \leq 2|z|^{m+1}, \quad z \in \mathbb{C}, |z| \leq 1/2.$$

Let $\mu(t)$, $t > 0$, be a probability measure on $(\mathbb{R}, \mathcal{B}(\mathbb{R}))$ given by

$$\mu(t)(A) = (1 - \bar{F}(t))^{-1} \mu(A \cap (-\infty, t]),$$

for any $A \in \mathcal{B}(\mathbb{R})$.

Let $\varphi(\cdot; \mu(t))$, $t > 0$, be the characteristic function of the probability measure $\mu(t)$, i.e.,

$$\varphi(\xi; \mu(t)) = \int_{\mathbb{R}} \exp(ix\xi) \mu(t)(dx), \quad \xi \in \mathbb{R}.$$

Proposition 6. *There is a $c_0 > 0$ such that for any $t \geq 2$, $\xi \in \mathbb{R}$ and integer n, m with $n \geq m$,*

$$|\varphi(n^{-1/2}\xi, \mu(t))|^n \leq (1 + \frac{c_0}{m} |\xi|^2)^{-m/4}.$$

Proposition 7. *Let ν be a probability measure on $(\mathbb{R}, \mathcal{B}(\mathbb{R}))$ such that $\int_{\mathbb{R}} x^2 \nu(dx) < \infty$. Also, assume that there is a constant $C > 0$ such that the characteristic function $\varphi(\cdot, \nu) : \mathbb{R} \rightarrow \mathbb{C}$ satisfies*

$$|\varphi(\xi; \nu)| \leq C(1 + |\xi|)^{-2}, \quad \xi \in \mathbb{R}.$$

Then for any $x \in \mathbb{R}$ and $v > 0$

$$\nu((x, \infty)) = \Phi_0(v^{-1/2}x) + \frac{1}{2\pi} \int_{\mathbb{R}} \frac{e^{-ix\xi}}{i\xi} (\varphi(\xi, \nu) - \exp(-\frac{v\xi^2}{2})) d\xi.$$

3 Estimate for moments

Let

$$\eta_k(t) = \int_{-\infty}^t x^k \mu(dx), \quad t > 0, \quad k = 1, 2,$$

and

$$\eta_3(t) = \int_1^t x^3 \mu(dx), \quad t > 1.$$

Then we see that

$$-\eta_1(t) = \int_t^\infty x \mu(dx) = \int_t^\infty \bar{F}(x) dx + t\bar{F}(t), \quad t > 0,$$

$$1 - \eta_2(t) = \int_t^\infty x^2 \mu(dx) = 2 \int_t^\infty x \bar{F}(x) dx + t^2 \bar{F}(t), \quad t > 0,$$

and

$$\eta_3(t) = \bar{F}(1) - t^3 \bar{F}(t) + 3 \int_1^t x^2 \bar{F}(x) dx \quad t > 1.$$

In particular, we see that

$$L(t) \leq 1 - \eta_2(t) \rightarrow 0, \quad t \rightarrow \infty, \quad (5)$$

$$\frac{1 - \eta_2(t)}{L(t)} \rightarrow \infty, \quad t \rightarrow \infty. \quad (6)$$

For any $\delta > 0$, let $t_n = n^{1/2} L(n^{1/2})^\delta$. Note that $n^{-1/2} t_n \rightarrow 0$, $n \rightarrow \infty$.

Proposition 8. *For any $\varepsilon > 0$, there is a $C > 0$ such that*

$$\frac{L(t_n)}{L(n^{1/2})} \leq CL(n^{1/2})^{-\varepsilon\delta} \quad (7)$$

$$n\bar{F}(t_n) \leq CL(n^{1/2})^{1-2\delta-\varepsilon\delta} \quad (8)$$

$$\eta_2(n^{1/2}) - \eta_2(t_n) \leq CL(n^{1/2})^{1-2\varepsilon\delta} \quad (9)$$

$$-n^{1/2}\eta_1(t_n) \leq CL(n^{1/2})^{1-2\delta} \quad (10)$$

$$n^{-1/2}\eta_3(t_n) \leq CL(n^{1/2}). \quad (11)$$

for any $n \geq 1$.

Proof. From Proposition 2, there is an $M(\varepsilon) > 0$ such that

$$\frac{L(t_n)}{L(n^{1/2})} = \frac{L(t_n)}{L(t_n L(n^{1/2})^{-\delta})} \leq M(\varepsilon) L(n^{1/2})^{-\varepsilon\delta}$$

Hence we have Inequality (7). Similarly, we see that

$$n\bar{F}(t_n) = L(n^{1/2})^{-2\delta} L(t_n) = L(n^{1/2})^{1-2\delta} \frac{L(t_n)}{L(n^{1/2})}$$

and

$$\begin{aligned} \eta_2(n^{1/2}) - \eta_2(t_n) &= L(t_n) - L(n^{1/2}) + 2 \int_{t_n}^{n^{1/2}} \frac{L(z)}{z} dz \\ &= L(t_n) - L(n^{1/2}) + 2L(t_n) \int_1^{L(t_n)^{-\delta}} \frac{L(t_n y)}{L(t_n) y} dy \\ &\leq L(t_n) - L(n^{1/2}) + 2L(t_n) M(\varepsilon) \int_1^{L(t_n)^{-\delta}} y^{-1+\varepsilon} dy \\ &\leq L(t_n) - L(n^{1/2}) + 2 \frac{M(\varepsilon)}{\varepsilon} L(t_n) (L(n^{1/2})^{-\varepsilon\delta} - 1) \end{aligned}$$

From Inequality (7), we have Inequalities (8) and (9).

Let

$$\varepsilon_1(t) = \frac{1}{t^{-1}L(t)} \int_t^\infty x^{-2}L(x)dx - 1$$

and

$$\varepsilon_3(t) = \frac{1}{tL(t)} \int_1^t L(x)dx - 1.$$

Then from Proposition 3 (1) and (2) we have $\varepsilon_1(t) \rightarrow 0$ and $\varepsilon_3(t) \rightarrow 0$ as $t \rightarrow \infty$.

Hence we see that

$$\begin{aligned} -n^{1/2}\eta_1(t_n) &= n^{1/2} \left(t_n \bar{F}(t_n) + \int_{t_n}^\infty \bar{F}(x)dx \right) = L(n^{1/2})^{-\delta} L(t_n) (2 + \varepsilon_1(t_n)) \\ &= (2 + \varepsilon_1(t_n)) L(n^{1/2})^{1-\delta} \frac{L(t_n)}{L(n^{1/2})} \end{aligned}$$

and

$$\begin{aligned} n^{-1/2}\eta_3(t_n) &= n^{-1/2} \bar{F}(1) + (2 + \varepsilon_3(t_n)) L(n^{1/2})^\delta L(t_n) \\ &= n^{-1/2} \bar{F}(1) + (2 + \varepsilon_3(t_n)) L(n^{1/2})^{1+\delta} \frac{L(t_n)}{L(n^{1/2})}. \end{aligned}$$

From Inequality (7), we have Inequalities (10) and (11). □

4 Asymptotic expansion of characteristic functions

Remind that $v_n = \int_{-\infty}^{n^{1/2}} x^2 \mu(dx)$ and $t_n = n^{1/2} L(n^{1/2})^\delta$.

In this section, we prove the following Lemma.

Lemma 1. *Let*

$$\begin{aligned} R_{n,0}(\xi) &= \exp\left(\frac{v_n}{2}\xi^2\right) \varphi\left(n^{-\frac{1}{2}}\xi; \mu(t_n)\right)^n - \left(1 + n\left(\varphi\left(n^{-\frac{1}{2}}\xi; \mu(t_n)\right) - 1\right) + \frac{v_n}{2}\xi^2\right), \\ R_{n,1}(\xi) &= \exp\left(\frac{v_n}{2}\xi^2\right) \varphi\left(n^{-\frac{1}{2}}\xi; \mu(t_n)\right)^n - 1, \\ R_{n,2}(\xi) &= \exp\left(\frac{v_n}{2}\xi^2\right) \varphi\left(n^{-\frac{1}{2}}\xi; \mu(t_n)\right)^{n-1} - 1. \end{aligned}$$

Then there is a $C > 0$ such that

$$|R_{n,0}(\xi)| \leq CL(n^{1/2})^{2-5\delta} |\xi|$$

and

$$|R_{n,1}(\xi)| + |R_{n,2}(\xi)| \leq CL(n^{1/2})^{1-2\delta}|\xi|$$

for any $n \geq 8$ and $\xi \in \mathbb{R}$ with $|\xi| \leq L(n^{1/2})^{-\delta}$.

As a corollary to Lemma 1, we have the following.

Corollary 1. *Let*

$$\begin{aligned} \tilde{R}_0(n, s) &= \mu(t_n)^{*n}((sn^{1/2}, \infty)) - \Phi_0(v_n^{-1/2}s) \\ &\quad - \frac{1}{2\pi} \int_{\mathbb{R}} \frac{e^{-is\xi}}{i\xi} \left(n(\varphi(n^{-\frac{1}{2}}\xi; \mu(t_n)) - 1) + \frac{v_n\xi^2}{2} \right) e^{-v_n\xi^2/2} d\xi \end{aligned}$$

$$\tilde{R}_{1,k}(n, s) = \mu(t_n)^{*(n-k)}((sn^{1/2}, \infty)) - \Phi_0(v_n^{-1/2}s), \quad k = 0, 1,$$

and

$$\tilde{R}_2(n, s) = \frac{1}{2\pi} \int_{\mathbb{R}} \left| \varphi(n^{-\frac{1}{2}}\xi; \mu(t_n))^{n-1} - e^{-\frac{v_n\xi^2}{2}} \right| d\xi.$$

Then there is a $C > 0$ such that for any $n \geq 1$, we have

$$|\tilde{R}_0(n, \xi)| \leq CL(n^{1/2})^{2-6\delta} \quad (12)$$

and

$$|\tilde{R}_{1,0}(n, \xi)| + |\tilde{R}_{1,1}(n, \xi)| + |\tilde{R}_2(n, \xi)| \leq CL(n^{1/2})^{1-4\delta}. \quad (13)$$

Proof. From Proposition 8, we see that

$$\begin{aligned} &\tilde{R}_0(n, s) \\ &= \frac{1}{2\pi} \int_{\mathbb{R}} \frac{e^{-is\xi}}{i\xi} \left(\varphi(n^{-\frac{1}{2}}\xi; \mu(t_n))^n - \left(1 + n(\varphi(n^{-\frac{1}{2}}\xi; \mu(t_n)) - 1) + \frac{v_n\xi^2}{2} \right) e^{-\frac{v_n\xi^2}{2}} \right) d\xi \\ &= \frac{1}{2\pi} \int_{\mathbb{R}} \frac{e^{-is\xi}}{i\xi} R_{n,0}(\xi) e^{-v_n\xi^2/2} d\xi. \end{aligned}$$

By Lemma 1, there is a $C_0 > 0$ such that

$$\int_{|\xi| \leq L(n^{1/2})^{-\delta}} \frac{|R_{n,0}(\xi)|}{|\xi|} d\xi \leq C_0 L(n^{1/2})^{2-6\delta}.$$

It is easy to see from Proposition 5 (1) that

$$n|\varphi(n^{-\frac{1}{2}}\xi; \mu(t_n)) - 1| \leq \frac{n^{1/2}|\eta_1(t_n)||\xi|}{1 - \bar{F}(t_n)} + \frac{|\xi|^2}{2(1 - \bar{F}(t_n))}, \quad \xi \in \mathbb{R}.$$

From the above inequality and Proposition 7, we see that for any $m \geq 2/\delta$, there is a $C_1 > 0$ such that for any $n \geq 2m$ and $\xi \in \mathbb{R}$ with $|\xi| \geq L(n^{1/2})^{-\delta}$,

$$|\varphi(n^{-\frac{1}{2}}\xi; \mu(t_n))| + \left| n(\varphi(n^{-\frac{1}{2}}\xi; \mu(t_n)) - 1) + 1 + \frac{v_n \xi^2}{2} \right| e^{-\frac{v_n \xi^2}{2}} \leq C_1 |\xi|^{-m}.$$

Hence we have

$$\begin{aligned} & \int_{|\xi| > L(n^{\frac{1}{2}})^{-\delta}} |\xi|^{-1} \left| \varphi(n^{-\frac{1}{2}}\xi; \mu(t_n))^n - \left(1 + n(\varphi(n^{-\frac{1}{2}}\xi; \mu(t_n)) - 1) + \frac{v_n \xi^2}{2} \right) e^{-\frac{v_n \xi^2}{2}} \right| d\xi \\ & \leq 2C_1 \int_{L(n^{1/2})^{-\delta}}^{\infty} |\xi|^{-m-1} d\xi = \frac{2C_1}{m} L(n^{1/2})^{m\delta} \leq \frac{2C_1}{m} L(n^{1/2})^2. \end{aligned}$$

Therefore we have Inequation (12).

We see also that

$$\begin{aligned} \tilde{R}_{1,k}(n, s) &= \frac{1}{2\pi} \int_{\mathbb{R}} \frac{e^{-is\xi}}{i\xi} \left(\varphi(n^{-\frac{1}{2}}\xi; \mu(t_n))^n - e^{-\frac{v_n \xi^2}{2}} \right) d\xi \\ &= \frac{1}{2\pi} \int_{\mathbb{R}} \frac{e^{-is\xi}}{i\xi} R_{n,1+k}(\xi) e^{-v_n \xi^2/2} d\xi \\ \tilde{R}_2(n, s) &= \frac{1}{2\pi} \int_{\mathbb{R}} |R_{n,2}(\xi)| e^{-v_n \xi^2/2} d\xi \end{aligned}$$

Similarly to the first equation, we have Inequation (13). □

We make some preparations to prove Lemma 1.

Let

$$R_0(n, \xi) = \varphi(n^{-1/2}\xi, \mu(t_n)) - \left(1 - v_n \frac{\xi^2}{2n} \right).$$

First we prove the following.

Proposition 9. *There is a constant $C > 0$ such that for any $n \geq 8$, and $\xi \in \mathbb{R}$ with $|\xi| \leq L(n^{1/2})^{-\delta}$,*

$$|nR_0(n, \xi)| \leq CL(n^{1/2})^{1-2\delta} |\xi|$$

and

$$n|\varphi(n^{-\frac{1}{2}}\xi; \mu(t_n)) - 1| \leq CL(n^{1/2})^{-\delta} |\xi|.$$

In particular,

$$\sup\{|nR_0(n, \xi)|; |\xi| \leq L(n^{1/2})^{-\delta}\} \rightarrow 0, \quad n \rightarrow \infty. \quad (14)$$

Proof. We can easily see that

$$\begin{aligned}\varphi(\xi; \mu(t)) &= \int_{\mathbb{R}} \exp(ix\xi) \mu(t)(dx) \\ &= 1 + \eta_1(t)(i\xi) + \eta_2(t) \frac{(i\xi)^2}{2} + \int_{-\infty}^1 r_{e,2}(\xi x) \mu(dx) \\ &\quad + \int_1^t r_{e,2}(\xi x) \mu(dx) + \frac{\bar{F}(t)}{1 - \bar{F}(t)} \int_{-\infty}^t r_{e,0}(\xi x) \mu(dx).\end{aligned}$$

Hence we have that

$$\begin{aligned}R_0(n, \xi) &= n^{-1/2} \eta_1(t_n)(i\xi) + (\eta_2(t_n) - \eta_2(n^{1/2})) \frac{(i\xi)^2}{2n} + \int_{-\infty}^1 r_{e,2}(n^{-1/2} \xi x) \mu(dx) \\ &\quad + \int_1^{t_n} r_{e,2}(n^{-1/2} \xi x) \mu(dx) + \frac{\bar{F}(t_n)}{1 - \bar{F}(t_n)} \int_{-\infty}^{t_n} r_{e,0}(n^{-1/2} \xi x) \mu(dx).\end{aligned}$$

Then we see that

$$\begin{aligned}n|R_0(n, \xi)| &\leq n^{\frac{1}{2}} |\eta_1(t_n)| |\xi| + (\eta_2(n^{\frac{1}{2}}) - \eta_2(t_n)) \frac{|\xi|^2}{2} + n^{-\frac{\delta_1}{2}} \int_{-\infty}^1 |x|^{2+\delta_0} \mu(dx) |\xi|^{2+\delta_0} \\ &\quad + \frac{1}{6} n^{-\frac{1}{2}} \eta_3(t_n) |\xi|^3 + 2n^{\frac{1}{2}} \bar{F}(t_n) \int_{\mathbb{R}} |x| \mu(dx) |\xi|, \quad \xi \in \mathbb{R}, t \geq 2,\end{aligned}$$

, where δ_0 is in (A2). Hence from Proposition 5, we see that there is a $C > 0$ such that

$$|nR_0(n, \xi)| \leq C (L(n^{1/2})^{1-2\delta} |\xi| + L(n^{1/2})^{1-\delta} |\xi|^2 + n^{-\delta_0/2} |\xi|^{2+\delta_0} + L(n^{1/2}) |\xi|^3).$$

Therefore we have the first inequality.

Since $n(\varphi(n^{-\frac{1}{2}} \xi; \mu(t_n)) - 1) = nR_0(n, \xi) - \eta_2(n^{1/2}) \xi^2 / 2$, we have the second inequality. \square

Let

$$R_{1,k}(n, \xi) = (n - k) \log \varphi(n^{-1/2} \xi; \mu(t_n)) - n(\varphi(n^{-\frac{1}{2}} \xi; \mu(t_n)) - 1), \quad k = 0, 1.$$

Proposition 10. *There is a $C > 0$ such that for any $\xi \in \mathbb{R}$ with $|\xi| \leq L(n^{1/2})^{-\delta}$,*

$$|R_{1,k}(n, \xi)| \leq C n^{-1} L(n^{1/2})^{-3\delta} |\xi|.$$

In particular

$$\sup\{|R_{1,k}(n, \xi)|; |\xi| \leq L(n^{1/2})^{-\delta}\} \rightarrow 0, \quad n \rightarrow \infty. \quad (15)$$

Proof. First, we have

$$\log \varphi(\xi, \mu(t)) = \varphi(\xi, \mu(t)) - 1 + r_{l,1}(\varphi(\xi, \mu(t)) - 1), \quad |\xi| \leq L(n^{1/2})^{-\delta}.$$

Hence we have

$$R_{1,k}(n, \xi) = -k \log \varphi(n^{-\frac{1}{2}}\xi; \mu(t_n)) + nr_{l,1}(\varphi(n^{-\frac{1}{2}}\xi; \mu(t_n)) - 1).$$

From Proposition 9, we see that there is a $C >$ such that

$$\begin{aligned} |R_{1,k}(n, \xi)| &\leq |\varphi(n^{-\frac{1}{2}}\xi; \mu(t_n)) - 1| + 2n|\varphi(n^{-\frac{1}{2}}\xi; \mu(t_n)) - 1|^2 \\ &\leq Cn^{-1}L(n^{1/2})^{-3\delta}|\xi|, \quad |\xi| \leq L(n^{1/2})^{-\delta}. \end{aligned}$$

□

Let us prove Lemma 1. Note that for $k = 0, 1$

$$\log(e^{v_n \xi^2/2} \varphi(n^{-\frac{1}{2}}\xi; \mu(t_n))^{n-k}) = nR_0(n, \xi) + R_{1,k}(n, \xi).$$

We see that

$$e^{v_n \xi^2/2} \varphi(n^{-\frac{1}{2}}\xi; \mu(t_n))^{n-k} = \exp(nR_0(n, \xi) + R_{1,k}(n, \xi))$$

Hence we see that

$$\begin{aligned} R_{n,0}(\xi) &= e^{v_n \xi^2/2} \varphi(n^{-\frac{1}{2}}\xi; \mu(t_n))^n - (1 + nR_0(n, \xi)) \\ &= \exp(nR_0(n, \xi)) - (1 + nR_0(n, \xi)) + \exp(nR_0(n, \xi))(\exp(R_{1,0}(n, \xi)) - 1) \end{aligned}$$

From Inequality (13), we see that there is a $C > 0$ such that

$$|R_{n,0}(\xi)| \leq C (|nR_0(n, \xi)|^2 + |R_{1,0}(n, \xi)|).$$

Therefore we have Inequality (12) from Proposition 9 and 10.

Proof of Inequality (13) is similar to Inequality (12).

5 Proof of Theorem 4

Note that

$$P\left(\sum_{l=1}^n X_l > sn^{1/2}\right) = \sum_{k=0}^n I_k(n, s)$$

where

$$I_k(n, s) = P\left(\sum_{l=1}^n X_l > sn^{1/2}, \sum_{l=1}^n 1_{\{X_l > t_n\}} = k\right), \quad k = 0, 1, \dots, n.$$

Then we have

$$I_k(n, s) = \binom{n}{k} P\left(\sum_{l=1}^n X_l > sn^{1/2}, X_i > t_n, i = 1, \dots, k, X_j \leq t_n, j = k+1, \dots, n\right),$$

for $k = 0, 1, \dots, n$.

Let $\bar{F}_{n,0}(x) = P(X_1 > n^{1/2}x, X_1 \leq t_n) = (1 - \bar{F}(t_n))\mu(t_n)((n^{1/2}x, \infty))$ and $\bar{F}_{n,1}(x) = P(X_1 > n^{1/2}x, X_1 > t_n)$. Note that $\bar{F}_{n,0}(x) + \bar{F}_{n,1}(x) = \bar{F}(n^{1/2}x)$.

Proposition 11. *There is a $C > 0$ such that*

$$\begin{aligned} & |I_0(n, s) - (1-n)\Phi_0(v_n^{-1/2}s) - \frac{1}{2}\Phi_2(v_n^{-1/2}s) - n \int_{\mathbb{R}} \bar{F}_{n,0}(s - v_n^{1/2}x)\Phi_1(x)dx| \\ & \leq CL(n^{1/2})^{2-5\delta}, \quad n \geq 1, s \geq 1. \end{aligned}$$

Proof. First, note that

$$\begin{aligned} & \int_{\mathbb{R}} \frac{\bar{F}_{n,0}(s - v_n^{1/2}x)}{1 - \bar{F}(t_n)}\Phi_1(x)dx - \Phi_0(v_n^{-1/2}s) \\ & = \int_s^\infty \frac{1}{2\pi} \left(\int_{\mathbb{R}} e^{-ix\xi}(\varphi(n^{-\frac{1}{2}}\xi; \mu(t_n)) - 1)e^{-\frac{v_n}{2}\xi^2} d\xi \right) dx \\ & = \frac{1}{2\pi} \int_{\mathbb{R}} \frac{e^{-is\xi}}{i\xi} (\varphi(n^{-\frac{1}{2}}\xi; \mu(t_n)) - 1)e^{-\frac{v_n}{2}\xi^2} d\xi. \end{aligned}$$

Hence we have

$$\begin{aligned} & n \left(\int_{\mathbb{R}} \frac{\bar{F}_{n,0}(s - v_n^{1/2}x)}{1 - \bar{F}(t_n)}\Phi_1(x)dx - \Phi_0(v_n^{-1/2}s) \right) + \frac{1}{2}\Phi_2(v_n^{-1/2}s) \\ & = \frac{1}{2\pi} \int_{\mathbb{R}} \frac{e^{-is\xi}}{i\xi} \left(n(\varphi(n^{-\frac{1}{2}}\xi; \mu(t_n)) - 1) + \frac{v_n\xi^2}{2} \right) e^{-v_n\xi^2/2} d\xi. \end{aligned}$$

By Corollary 1, we see that there is a $C > 0$ such that for any $n \geq 1$, we have

$$\begin{aligned} & |\mu(t_n)^{*n}((sn^{1/2}, \infty)) - (1-n)\Phi_0(v_n^{-1/2}s) - \frac{v_n^{-1/2}}{2}\Phi_2(s) \\ & \quad - n \int_{\mathbb{R}} \frac{\bar{F}_{n,0}(s - v_n^{1/2}x)}{1 - \bar{F}(t_n)}\Phi_1(x)dx| \\ & \leq CL(n^{1/2})^{2-5\delta}. \end{aligned}$$

We decompose $I_0(n, s)$ into three parts, i.e.,

$$\begin{aligned} I_0(n, s) &= (1 - n\bar{F}(t_n))^n \mu(t_n)^{*n}((sn^{1/2}, \infty)) \\ &= I_{0,0}(n, s) + I_{0,1}(n, s) + I_{0,2}(n, s), \end{aligned}$$

where

$$\begin{aligned} I_{0,0}(n, s) &= (1 - \bar{F}(t_n)) \mu(t_n)^{*n}((sn^{1/2}, \infty)), \\ I_{0,1}(n, s) &= -n\bar{F}(t_n) \mu(t_n)^{*n}((sn^{1/2}, \infty)), \\ I_{0,2}(n, s) &= (1 - n\bar{F}(t_n))^n - 1 + (n+1)\bar{F}(t_n) \mu(t_n)^{*n}((sn^{1/2}, \infty)). \end{aligned}$$

Since

$$\begin{aligned} &(1 - \bar{F}(t_n)) \left((1 - n)\Phi_0(v_n^{-1/2}s) + \frac{\Phi_2(v_n^{-1/2}s)}{2} + n \int_{\mathbb{R}} \frac{\bar{F}_{n,0}(s - v_n^{1/2}x)}{1 - \bar{F}(t_n)} \Phi_1(x) dx \right) \\ &= (1 - n + n\bar{F}(t_n))\Phi_0(v_n^{-1/2}s) + \frac{\Phi_2(v_n^{-1/2}s)}{2} + n \int_{\mathbb{R}} \bar{F}_{n,0}(s - v_n^{1/2}x) \Phi_1(x) dx \\ &\quad - \bar{F}(t_n)\Phi_0(v_n^{-1/2}s) - \bar{F}(t_n) \frac{\Phi_2(v_n^{-1/2}s)}{2}, \end{aligned}$$

we have

$$\begin{aligned} &|I_{0,0}(n, s) - (1 - n + n\bar{F}(t_n))\Phi_0(v_n^{-1/2}s) - \frac{\Phi_2(v_n^{-1/2}s)}{2} - n \int_{\mathbb{R}} \bar{F}_{n,0}(s - v_n^{1/2}x) \Phi_1(x) dx| \\ &\leq CL(n^{1/2})^{2-5\delta}. \end{aligned}$$

By Corollary 1, we see that

$$|I_{0,1}(n, s) + n\bar{F}(t_n)\Phi_0(v_n^{-1/2}s)| \leq CL(n^{1/2})^{2-5\delta}.$$

Note that $|(1-x)^n - (1-nx)| \leq n^2x^2$ for any $x \in [0, 1]$, $n \geq 1$. Hence we have

$$|I_{0,2}(n, s)| \leq Cn^2\bar{F}(t_n)^2 \leq CL(n^{1/2})^{2-5\delta}$$

Therefore we have our assertion. □

Proposition 12. *There is a $C > 0$ such that*

$$|I_1(n, s) - n \int_{\mathbb{R}} \bar{F}_{n,1}(s - v_n^{1/2}x) \Phi_1(x) dx| \leq CL(n^{1/2})^{2-6\delta}, \quad n \geq 1, s \geq 1.$$

Proof.

$$\begin{aligned}
I_1(n, s) &= n(1 - \bar{F}(t_n))^{n-1} \int_{\mathbb{R}} P(X_1 + x > sn^{1/2}, X_1 > t_n) \mu(t_n)^{* (n-1)}(dx) \\
&= n(1 - \bar{F}(t_n))^{n-1} \int_{\mathbb{R}} \bar{F}((sn^{1/2} - x) \vee t_n) \mu(t_n)^{* (n-1)}(dx) \\
&= nJ_0(n, s) + nJ_1(n, s) + nJ_2(n, s),
\end{aligned}$$

where

$$\begin{aligned}
J_0(n, s) &= \int_{-\infty}^{sn^{1/2} - t_n} \bar{F}(sn^{1/2} - x) \mu(t_n)^{* (n-1)}(dx), \\
J_1(n, s) &= \bar{F}(t_n) \int_{sn^{1/2} - t_n}^{\infty} \mu(t_n)^{* (n-1)}(dx) = \bar{F}(t_n) \mu(t_n)^{* (n-1)}((s - L(n^{1/2})^\delta) n^{1/2}, \infty),
\end{aligned}$$

and

$$J_2(n, s) = ((1 - \bar{F}(t_n))^{n-1} - 1) \int_{\mathbb{R}} \bar{F}((sn^{1/2} - x) \vee t_n) \mu(t_n)^{* (n-1)}(dx).$$

We see that

$$\begin{aligned}
&J_0(n, s) \\
&= \int_{-\infty}^{sn^{1/2} - t_n} \bar{F}(sn^{1/2} - x) \mu(t_n)^{* (n-1)}(dx) \\
&= \int_{-\infty}^{sn^{1/2} - t_n} \bar{F}(sn^{1/2} - x) \left(\frac{1}{2\pi} \int_{\mathbb{R}} e^{-in^{-1/2}x\xi} \left(\varphi(n^{-\frac{1}{2}}\xi; \mu(t_n))^{n-1} - e^{-\frac{v_n \xi^2}{2}} \right) \right. \\
&\quad \left. + v_n^{-1/2} \Phi_1(n^{-1/2} v_n^{-1/2} x) \right) n^{-1/2} dx \\
&= \int_{-\infty}^{s - L(n^{1/2})^\delta} \bar{F}((s - x)n^{1/2}) \frac{1}{2\pi} \left(\int_{\mathbb{R}} e^{-ix\xi} (\varphi(n^{-\frac{1}{2}}\xi; \mu(t_n))^{n-1} - e^{-\frac{v_n \xi^2}{2}}) d\xi \right) dx \\
&\quad + \int_{-\infty}^{v_n^{-1/2}(s - L(n^{1/2})^\delta)} \bar{F}((s - v_n^{1/2}x)n^{1/2}) \Phi_1(x) dx
\end{aligned}$$

Hence from Corollary 1, we see that there is a $C > 0$ such that

$$\begin{aligned}
&n |J_0(n, s) - \int_{-\infty}^{v_n^{-1/2}(s - L(n^{1/2})^\delta)} \bar{F}((s - v_n^{1/2}x)n^{1/2}) \Phi_1(x) dx| \\
&\leq CL(n^{1/2})^{1-4\delta} n^{1/2} \int_{t_n}^{\infty} \bar{F}(x) dx \\
&\leq CL(n^{1/2})^{1-4\delta} n^{1/2} |\eta_1(t_n)|
\end{aligned}$$

We also see that

$$\begin{aligned}
&n |J_1(n, s) - \bar{F}(t_n) \Phi_0(v_n^{-1/2}(s - L(n^{1/2})^\delta))| \\
&\leq Cn \bar{F}(t_n) |\mu(t_n)^{* (n-1)}((s - L(n^{1/2})^\delta) n^{1/2}, \infty) - \Phi_0(v_n^{-1/2}(s - L(n^{1/2})^\delta))| \\
&\leq Cn \bar{F}(t_n) L(n^{1/2})^{1-4\delta}
\end{aligned}$$

and

$$|nJ_2(n, s)| \leq (n\bar{F}(t_n))^2 \leq L(n^{1/2})^{2-6\delta}.$$

Note that

$$\begin{aligned} & \int_{-\infty}^{v_n^{-1/2}(s-L(n^{1/2})^\delta)} \bar{F}((s-v_n^{1/2}x)n^{1/2})\Phi_1(x)dx + \bar{F}(t_n)\Phi_0(v_n^{-1/2}(s-L(n^{1/2})^\delta)) \\ &= \int_{\mathbb{R}} \bar{F}_{n,1}(s-v_n^{1/2}x)\Phi_1(x)dx. \end{aligned}$$

Therefore we have our assertion. □

Let us prove Theorem 4.

From Proposition 11 and 12, we see that there is a $C > 0$ such that

$$\begin{aligned} & |I_0(n, s) + I_1(n, s) \\ & \quad - (1-n)\Phi_0(v_n^{-1/2}s) - \frac{1}{2}\Phi_2(v_n^{-1/2}s) - n \int_{\mathbb{R}} \bar{F}(n^{1/2}(s-v_n^{1/2}x))\Phi_1(x)dx| \\ & \leq CL(n^{1/2})^{2-6\delta} \end{aligned}$$

Note that

$$\begin{aligned} & \int_{\mathbb{R}} \bar{F}(n^{\frac{1}{2}}(s-v_n^{1/2}x))\Phi_1(x)dx - \Phi_0(v_n^{-1/2}s) \\ &= \int_{-\infty}^{v_n^{-1/2}s} \bar{F}(n^{\frac{1}{2}}(s-v_n^{1/2}x))\Phi_1(x)dx \\ & \quad + \int_{v_n^{-1/2}s}^{\infty} (\bar{F}(n^{\frac{1}{2}}(s-v_n^{1/2}x)) - 1_{\{v_n^{1/2}x>s\}})\Phi_1(x)dx \\ &= \int_{-\infty}^{v_n^{-1/2}s} \bar{F}(n^{\frac{1}{2}}v_n^{1/2}(v_n^{-\frac{1}{2}}s-x))\Phi_1(x)dx - \int_{v_n^{-1/2}s}^{\infty} F(n^{\frac{1}{2}}(s-v_n^{1/2}x))\Phi_1(x)dx \end{aligned}$$

and

$$n \int_{v_n^{-1/2}s}^{\infty} F(n^{1/2}(s-v_nx))\Phi_1(x)dx = n^{1/2} \int_{-\infty}^0 F(y)\Phi_1(v_n^{-1/2}s - n^{-1/2}v_n^{-1/2}y)v_n^{-1/2}dy.$$

Let $R(z, y) = \Phi_1(z-y) - \Phi_1(z) - \Phi_2(z)y$, for $z > 0, y \leq 0$, then we see that there is a $C_1 > 0$ such that

$$|R(s, y)| \leq C_1|y|^{1+\delta_0}.$$

Hence we have

$$\begin{aligned}
& n \left| \int_{v_n^{-1/2}s}^{\infty} F(n^{1/2}(s - v_n x)) \Phi_1(x) dx - \sum_{k=1}^2 v_n^{-k/2} n^{-k/2} \Phi_k(v_n^{-1/2}s) \int_{-\infty}^0 y^{k-1} F(y) dy \right| \\
&= \left| n^{1/2} \int_{-\infty}^0 R(v_n^{-1/2}s, n^{-1/2}v_n^{-1/2}y) F(y) dy \right| \\
&\leq C_1 n^{-\delta_1/2} v_n^{-(1+\delta_1)/2} \int_{-\infty}^0 |y|^{1+\delta_0} F(y) dy \\
&\leq C n^{-\delta_1/2},
\end{aligned}$$

where $C = C_1 v_1^{-(1+\delta_1)/2} \int_{-\infty}^0 y^{1+\delta_1} F(y) dy < \infty$.

Since

$$\int_{-\infty}^0 F(y) dy = \int_{-\infty}^0 y \mu(dy) = - \int_0^{\infty} y \mu(dy)$$

and

$$- \int_{-\infty}^0 y F(y) dy = \frac{1}{2} \int_{-\infty}^0 y^2 \mu(dy) = \frac{v_n}{2} - \frac{1}{2} \int_0^{n^{1/2}} y^2 \mu(dy),$$

we see that

$$\frac{1}{2} \Phi_2(v_n^{-1/2}s) + v_n^{-1} \Phi_2(v_n^{-1/2}s) \int_{-\infty}^0 y^2 F(y) dy = v_n^{-1} \frac{\Phi_2(v_n^{-1/2}s)}{2} \int_0^{n^{1/2}} y^2 \mu(dy)$$

Therefore we have

$$\begin{aligned}
& \left| (1-n) \Phi_0(v_n^{-1/2}s) + \frac{1}{2} \Phi_2(v_n^{-1/2}s) + n \int_{\mathbb{R}} \bar{F}(n^{1/2}(s - v_n^{1/2}x)) \Phi_1(x) dx - H(n, v_n^{-1/2}s) \right| \\
&\leq C n^{-\delta_0/2}.
\end{aligned}$$

We also see that

$$\begin{aligned}
\sum_{k=2}^n I_k(n, s) &\leq \sum_{k=2}^n \frac{n(n-1)}{k(k-1)} \binom{n-2}{k-2} \bar{F}(t_n)^k (1 - \bar{F}(t_n))^{n-k} \leq \frac{n(n-1)}{2} \bar{F}(t_n)^2 \\
&\leq L(n^{1/2})^{2-5\delta}.
\end{aligned}$$

This completes the proof of Theorem 4.

6 Some Estimations

Let

$$\begin{aligned}\hat{F}_n(s) &= \int_{-\infty}^s \bar{F}((s-x)v_n^{1/2}n^{1/2})\Phi_1(x)dx, \\ A(n, s) &= n\hat{F}_n(s) - v_n^{-1/2}n^{1/2}\Phi_1(s) \int_0^\infty x\mu(dx) - \frac{v_n^{-1}}{2}\Phi_2(s) \int_0^{n^{1/2}} x^2\mu(dx), \\ &= n\hat{F}_n(s) - v_n^{-1/2}n^{1/2}\Phi_1(s) \int_0^\infty \bar{F}(x)dx \\ &\quad - v_n^{-1}\Phi_2(s) \left(\int_0^{n^{1/2}} x\bar{F}(x)dx - \frac{L(n^{1/2})}{2} \right).\end{aligned}$$

Then we have

$$H(n, s) = \Phi_0(s) + A(n, s).$$

Let

$$H_0(n, s) = \Phi_0(s) + n\bar{F}(v_n^{1/2}n^{1/2}s).$$

In this section we prove the following Lemma.

Lemma 2.

$$\sup_{s \in [1, \infty)} \left| \frac{H(n, s)}{H_0(n, s)} - 1 \right| \rightarrow 0, \quad n \rightarrow \infty.$$

Let $u_n = v_n^{1/2}n^{1/2}$, $\alpha_n = L(u_n)^{1/3}$ and $\beta_n = L(u_n)^{-1/12}$.

Proposition 13. *For any $\varepsilon > 0$, there is a $C > 0$ such that*

$$\frac{1}{nF(u_ns)} \leq CL(u_n)^{-1}s^{2+\varepsilon}, \quad s \in [1, \infty).$$

In particular, for $s > \beta_n$ we have

$$\frac{1}{nF(u_ns)} \leq Cs^{14+\varepsilon}.$$

Proof. From Proposition 8 we see that for any $\varepsilon > 0$ there is a $C > 0$ such that

$$\begin{aligned}\frac{1}{nF(u_ns)} &= v_ns^2 \frac{1}{L(u_n)} \frac{L(u_n)}{L(u_ns)} \\ &\leq CL(u_n)^{-1}s^{2+\varepsilon}\end{aligned}$$

Since $L(u_n)^{-1} = \beta_n^{12} \leq s^{12}$ for $s > \beta_n$, we have the second inequality. \square

Let $n\hat{F}_n(s) = \sum_{k=1}^4 I_k(n, s)$, where

$$\begin{aligned} I_1(n, s) &= n \int_{s-\alpha_n}^s \bar{F}((s-x)u_n) \Phi_1(x) dx \\ I_2(n, s) &= n \int_{\sqrt{7/8}s}^{s-\alpha_n} \bar{F}((s-x)u_n) \Phi_1(x) dx \\ I_3(n, s) &= n \int_{-s}^{\sqrt{7/8}s} \bar{F}((s-x)u_n) \Phi_1(x) dx \\ I_4(n, s) &= n \int_{-\infty}^{-s} \bar{F}((s-x)u_n) \Phi_1(x) dx \end{aligned}$$

Let

$$R(n, s, y) = \Phi_1(s - u_n^{-1}y) - (\Phi_1(s) + u_n^{-1}y\Phi_2(s)), \quad \text{for } n \geq 1, s, y \in [1, \infty).$$

Proposition 14.

$$\sup_{s \in [1, \infty)} H_0(n, s)^{-1} |I_1(n, s) - \sum_{k=1}^2 v_n^{-k/2} n^{-(k-2)/2} \Phi_k(s) \int_0^{\alpha_n u_n} y^{k-1} \bar{F}(y) dy| \rightarrow 0, \quad n \rightarrow \infty.$$

Proof. We see that

$$\begin{aligned} & I_1(n, s) - \sum_{k=1}^2 v_n^{-k/2} n^{-(k-2)/2} \Phi_k(s) \int_0^{\alpha_n u_n} y^{k-1} \bar{F}(y) dy \\ &= nu_n^{-1} \int_0^{\alpha_n u_n} \bar{F}(y) (\Phi_1(s - u_n^{-1}y) - \Phi_1(s) - u_n^{-1}y\Phi_2(s)) dy \\ &= nu_n^{-1} \int_0^{\alpha_n u_n} \bar{F}(y) R(n, s, y) dy \end{aligned}$$

Note that for any $y \in [0, \alpha_n u_n]$,

$$\begin{aligned} |R(n, s, y)| &\leq u_n^{-2} y^2 \sup_{z \in [s-\alpha_n, s]} |\Phi_3(z)| \\ &\leq C_0 n^{-1} y^2 (1+s)^2 \Phi_1(s - \alpha_n) \\ &\leq C_0^2 n^{-1} y^2 (1+s)^3 \Phi_0(s) \exp(\alpha_n s) \end{aligned}$$

Hence for all $s \in [1, \infty)$

$$\begin{aligned} & |I_1(n, s) - \sum_{k=1}^2 v_n^{-k/2} n^{-(k-2)/2} \Phi_k(s) \int_0^{\alpha_n u_n} y^{k-1} \bar{F}(y) dy| \\ &\leq 8C_0 \sup\{z^2 \bar{F}(z); z \geq 0\} \alpha_n s^3 \Phi_0(s) \exp(\alpha_n s) \end{aligned}$$

Since $\alpha_n \beta_n^3 = L(u_n)^{1/12} \rightarrow 0$, $n \rightarrow \infty$, we have

$$\sup_{s \leq \beta_n} \Phi_0(s)^{-1} |I_1(n, s) - \sum_{k=1}^2 v_n^{-k/2} n^{-(k-2)/2} \Phi_k(s) \int_0^{\alpha_n u_n} y^{k-1} \bar{F}(y) dy| \rightarrow 0, \quad n \rightarrow \infty.$$

From Proposition 13, we see that for any $\varepsilon > 0$ there is a $C(\varepsilon) > 0$ such that

$$(n\bar{F}(u_n s))^{-1} \leq C(\varepsilon) s^{14+\varepsilon}.$$

Hence we see that for $s > \beta_n$,

$$\begin{aligned} & (n\bar{F}(u_n s))^{-1} |I_1(n, s) - \sum_{k=1}^2 v_n^{-k/2} n^{-(k-2)/2} \Phi_k(s) \int_0^{\alpha_n u_n} y^{k-1} \bar{F}(y) dy| \\ & \leq 8C(\varepsilon) C_0^2 \sup\{z^2 \bar{F}(z); z \geq 0\} \alpha_n s^{17+\varepsilon} \Phi_0(s) \exp(\alpha_n s). \end{aligned}$$

Since $\sup_{n \geq 1} \sup_{s > \beta_n} s^{17+\varepsilon} \Phi_0(s) \exp(\alpha_n s) < \infty$, we have

$$\sup_{s > \beta_n} (n\bar{F}(u_n s))^{-1} |I_1(n, s) - \sum_{k=1}^2 v_n^{-k/2} n^{-(k-2)/2} \Phi_k(s) \int_0^{\alpha_n u_n} y^{k-1} \bar{F}(y) dy| \rightarrow 0, \quad n \rightarrow \infty.$$

Therefore we have our assertion. \square

Proposition 15.

$$\sup_{s \in [1, \infty)} H_0(n, s)^{-1} |I_2(n, s) - \sum_{k=1}^2 v_n^{-k/2} n^{(2-k)/2} \Phi_k(s) \int_{\alpha_n u_n}^{(1-\sqrt{7/8})u_n s} y^{k-1} \bar{F}(y) dy| \rightarrow 0, \quad n \rightarrow \infty.$$

Proof. Similarly to Proposition 14, we see that

$$\begin{aligned} & |I_2(n, s) - \sum_{k=1}^2 v_n^{-k/2} n^{(2-k)/2} \Phi_k(s) \int_{\alpha_n u_n}^{(1-\sqrt{7/8})u_n s} y^{k-1} \bar{F}(y) dy| \\ & \leq n u_n^{-1} \int_{\alpha_n u_n}^{(1-\sqrt{7/8})u_n s} \bar{F}(y) |R(n, s, y)| dy \\ & \leq n u_n^{-3} \bar{F}(u_n \alpha_n) C_0 (1+s)^2 \left(\sup_{z \in [\sqrt{7/8}s, s]} |\Phi_1(z)| \right) \int_{u_n \alpha_n}^{(1-\sqrt{7/8})u_n s} y^2 dy \\ & \leq 4C_0 n \bar{F}(u_n \alpha_n) s^5 \Phi_1(\sqrt{7/8}s) \\ & \leq 4C_0^{1+7/8} n \bar{F}(u_n \alpha_n) s^6 \Phi_0(s)^{7/8}. \end{aligned}$$

Since $H_0(n, s)^{-1} \leq \Phi_0(s)^{-6/7} (n\bar{F}(u_n s))^{-1/7}$, it is easy to see that for any $\varepsilon \in (0, 4/7)$, there is a $C_1 > 0$ such that

$$\begin{aligned} & H_0(n, s)^{-1} |I_2(n, s) - \sum_{k=1}^2 v_n^{-k/2} n^{(2-k)/2} \Phi_k(s) \int_{\alpha_n u_n}^{(1-\sqrt{7/8})u_n s} y^{k-1} \bar{F}(y) dy| \\ & \leq C_1 s^{6+2/7+\varepsilon} \Phi_0(s)^{7/8-6/7} L(u_n)^{(1-\varepsilon)/3-1/7}. \end{aligned}$$

Since $\sup_{s \geq 1} \{s^{6+2/7+\varepsilon} \Phi_0(s)^{7/8-6/7}\} < \infty$ and $(1-\varepsilon)/3 - 1/7 > 0$, we have

$$\sup_{s \geq 1} H_0(n, s)^{-1} |I_2(n, s) - \sum_{k=1}^2 v_n^{-k/2} n^{(2-k)/2} \Phi_k(s) \int_{\alpha_n u_n}^{(1-\sqrt{7/8})u_n s} y^{k-1} \bar{F}(y) dy| \rightarrow 0, \quad n \rightarrow \infty.$$

□

Proposition 16.

$$\sup_{s \in [1, \infty)} H_0(n, s)^{-1} |I_3(n, s) - n\bar{F}(u_n s)| \rightarrow 0, \quad n \rightarrow \infty.$$

Proof.

$$\begin{aligned} I_3(n, s) &= n\bar{F}(u_n s) \int_{-s}^{\sqrt{7/8}s} \frac{\bar{F}(u_n(s-x))}{\bar{F}(u_n s)} \Phi_1(x) dx \\ &= n\bar{F}(u_n s) \int_{-s}^{\sqrt{7/8}s} \left(1 - \frac{x}{s}\right)^{-2} \frac{L(u_n(s-x))}{L(u_n s)} \Phi_1(x) dx. \end{aligned}$$

It is easy to see that there is a $C_1 > 0$ such that

$$\begin{aligned} \Phi_0(s)^{-1} &\leq C_1 L(u_n)^{-2/3}, \quad n \geq 1, s \in [1, (-\log L(u_n))^{1/2}], \\ \left| \int_{-s}^{\sqrt{7/8}s} \frac{\bar{F}(u_n(s-x))}{\bar{F}(u_n s)} \Phi_1(x) dx \right| &\leq C_1, \quad n \geq 1, s \in [1, \infty). \end{aligned}$$

Then we have

$$\begin{aligned} &\sup_{s \leq (-\log L(u_n))^{1/2}} H_0(n, s)^{-1} |I_3(n, s) - n\bar{F}(u_n s)| \\ &\leq C_1(C_1 + 1) L(u_n)^{-2/3} n\bar{F}(u_n) \\ &\leq C_1(C_1 + 1) v_n^{-1} L(u_n)^{1/3} \rightarrow 0, \quad n \rightarrow \infty. \end{aligned}$$

We take $M > 1$ arbitrarily, then $(-\log L(u_n))^{1/4} > M$ for sufficiently large n .

Hence we see that for $s > (-\log L(u_n))^{1/2}$

$$\begin{aligned}
& \left| \int_{-s}^{\sqrt{7/8}s} \left(1 - \frac{x}{s}\right)^{-2} \frac{L(u_n(s-x))}{L(u_ns)} \Phi_1(x) dx - 1 \right| \\
& \leq \left| \int_{-s}^{\sqrt{7/8}s} \left\{ \left(1 - \frac{x}{s}\right)^{-2} - 1 \right\} \frac{L(u_n(s-x))}{L(u_ns)} \Phi_1(x) dx \right| \\
& \quad + \left| \int_{-s}^{\sqrt{7/8}s} \left(\frac{L(u_n(s-x))}{L(u_ns)} - 1 \right) \Phi_1(x) dx \right| + \int_{[-s, \sqrt{7/8}s]^c} \Phi_1(x) dx \\
& \leq 2 \left(\int_{-M}^M \left| \left(1 - \frac{x}{s}\right)^{-2} - 1 \right| \Phi_1(x) dx + 8\Phi_0(M) \right) \\
& \quad + \sup_{t > (-\log L(u_n))^{1/2}} \sup_{1 - \sqrt{7/8} \leq a \leq 1} \left| \frac{L(at)}{L(t)} - 1 \right| + 2\Phi_0(\sqrt{7/8}s)
\end{aligned}$$

Hence we have

$$\sup_{s > (-\log L(u_n))^{1/2}} |n\bar{F}(u_ns)|^{-1} |I_3(n, s) - n\bar{F}(u_ns)| \rightarrow 0, \quad n \rightarrow \infty.$$

So we have our assertion. \square

Proposition 17.

$$\sup_{s \in [1, \infty)} \frac{I_4(n, s)}{H_0(n, s)} \rightarrow 0, \quad n \rightarrow \infty.$$

Proof. $|I_4(n, s)| \leq n\bar{F}(2u_ns)\Phi_0(s)$. Hence we have

$$\Phi_0(s)^{-1} |I_4(n, s)| \leq n\bar{F}(2u_ns) \leq n\bar{F}(u_n) \rightarrow 0, \quad n \rightarrow \infty.$$

\square

Proposition 18.

$$\sup_{s \in [1, \infty)} H_0(n, s)^{-1} |v_n^{-1/2} n^{1/2} \Phi_1(s) \int_{\sqrt{7/8}u_ns}^{\infty} \bar{F}(y) dy| \rightarrow 0, \quad n \rightarrow \infty,$$

and

$$\sup_{s \in [1, \infty)} H_0(n, s)^{-1} |v_n^{-1} \Phi_2(s) \left(\int_{\sqrt{7/8}u_ns}^{n^{1/2}} y \bar{F}(y) dy + L(n^{1/2}) \right)| \rightarrow 0, \quad n \rightarrow \infty.$$

Proof. From Proposition 3 (2), we see that there is a $C_1 > 0$ such that

$$n^{1/2} \int_{(1-\sqrt{7/8})u_n s}^{\infty} \bar{F}(y) dy \leq C_1 s^{-1} L((1 - \sqrt{7/8})u_n s).$$

We can easily see that

$$\sup_{s \in [1, \beta_n]} \Phi_0(s)^{-1} n^{1/2} \Phi_1(s) \int_{(1-\sqrt{7/8})n^{1/2}s}^{\infty} \bar{F}(y) dy \rightarrow 0, \quad n \rightarrow \infty,$$

and

$$\sup_{s \in [\beta_n, \infty)} (n\bar{F}(n^{1/2}s))^{-1} n^{1/2} \Phi_1(s) \int_{(1-\sqrt{7/8})n^{1/2}s}^{\infty} \bar{F}(y) dy \rightarrow 0, \quad n \rightarrow \infty.$$

Also we see that for any $\varepsilon \in (0, 1)$, there is a $C_2 > 0$ such that

$$\int_{(1-\sqrt{7/8})u_n s}^{n^{1/2}} y \bar{F}(y) dy = \int_{(1-\sqrt{7/8})v_n^{1/2}s}^1 \frac{L(n^{1/2}x)}{x} dx \leq C_2 L(n^{1/2}) s^\varepsilon.$$

Hence we can easily see that

$$\sup_{s \in [1, \beta_n]} \Phi_0(s)^{-1} |\Phi_2(s) \left(\int_{(1-\sqrt{7/8})u_n s}^{n^{1/2}} y \bar{F}(y) dy + L(n^{1/2}) \right)| \rightarrow 0, \quad n \rightarrow \infty,$$

and

$$\sup_{s \in [\beta_n, \infty)} (n\bar{F}(n^{1/2}s))^{-1} |\Phi_2(s) \left(\int_{(1-\sqrt{7/8})u_n s}^{n^{1/2}} y \bar{F}(y) dy + L(n^{1/2}) \right)| \rightarrow 0, \quad n \rightarrow \infty.$$

Therefore we have our assertion. \square

Now let us prove Lemma 2. Note that $H(n, s) - H_0(n, s) = A(n, s) - n\bar{F}(sn^{1/2})$.

So Proposition 14, 15, 16, 17 and 18 imply Lemma 2.

7 Proof of Theorem 2

First we prove the following Lemma.

Lemma 3. *For any $\beta > 0$ and $\delta \in (0, 1)$, there is a $C > 0$ such that*

$$\sup_{s > L(n^{1/2})^{-\beta}} \left| \frac{P(\sum_{k=1}^n X_k > sn^{1/2})}{H(n, v_n^{-1/2}s)} - 1 \right| \leq CL(n^{1/2})^{1-\delta}$$

We make some preparation to prove Lemma 3. Similarly to Proposition 26 in [2], we can prove the following.

Proposition 19. (1) For any $t, s > 0$, and $n \geq 2$,

$$P\left(\sum_{k=2}^n X_k 1_{\{X_k \leq tn^{1/2}\}} > sn^{\frac{1}{2}}\right) \leq \exp\left(\frac{6}{t^2} - \frac{s}{t}\right).$$

(2) For any $s, t > 0$, $\varepsilon \in (0, 1)$ with $t < (1 - \varepsilon)s$,

$$\begin{aligned} & \left| P\left(\sum_{k=1}^n X_k > sn^{\frac{1}{2}}\right) - nP\left(X_1 + \sum_{k=2}^n X_k 1_{\{X_k \leq tn^{1/2}\}} > sn^{\frac{1}{2}}, \sum_{k=2}^n X_k 1_{\{X_k \leq tn^{1/2}\}} \leq \varepsilon sn^{\frac{1}{2}}\right) \right| \\ & \leq 2n(n-1)\bar{F}(tn^{\frac{1}{2}})^2 + \exp\left(\frac{6}{t^2} - \frac{s}{t}\right) + n\bar{F}(tn^{\frac{1}{2}})\exp\left(\frac{6}{t^2} - \frac{\varepsilon s}{2t}\right). \end{aligned}$$

Also we prove the following for the proof of Lemma 3.

Proposition 20. For any $\gamma, \delta, \varepsilon \in (0, 1)$ and $\beta > 0$, there is a $C > 0$ such that

$$\begin{aligned} & \left| P\left(X_1 + \sum_{k=2}^n X_k 1_{\{X_k \leq s\gamma n^{1/2}\}} > sn^{1/2}, \sum_{k=2}^n X_k 1_{\{X_k \leq s\gamma n^{1/2}\}} \leq \varepsilon sn^{1/2}\right) \right. \\ & \quad \left. - \int_{-\infty}^{\varepsilon v_n^{-1/2} s} \bar{F}(n^{1/2}(s - v_n^{1/2}x))\Phi_1(x)dx \right| \\ & \leq C\bar{F}((1 - \varepsilon)n^{1/2}s)L(n^{1/2})^{1-3\delta}, \quad \text{for } s > L(n^{1/2})^{-\beta}. \end{aligned}$$

Proof. Let $t_n = L(n^{1/2})^\delta n^{1/2}$.

It is easy to see that there is a $C_1 > 0$ such that for $s > L(n^{1/2})^{-\beta}$

$$\begin{aligned} & \left| P\left(X_1 + \sum_{k=2}^n X_k 1_{\{X_k \leq s\gamma n^{1/2}\}} > sn^{\frac{1}{2}}, \sum_{k=2}^n X_k 1_{\{X_k \leq s\gamma n^{1/2}\}} \leq \varepsilon sn^{\frac{1}{2}}\right) \right. \\ & \quad \left. - P\left(X_1 + \sum_{k=2}^n X_k > sn^{\frac{1}{2}}, \sum_{k=2}^n X_k \leq \varepsilon sn^{\frac{1}{2}}, X_2 \leq t_n, \dots, X_n \leq t_n\right) \right| \\ & \leq C_1\bar{F}((1 - \varepsilon)n^{\frac{1}{2}}s)L(n^{\frac{1}{2}})^{1-3\delta}. \end{aligned}$$

We also see that

$$\begin{aligned} & P\left(X_1 + \sum_{k=2}^n X_k > sn^{1/2}, \sum_{k=2}^n X_k \leq \varepsilon sn^{1/2}, X_2 \leq t_n, \dots, X_n \leq t_n\right) \\ & = (1 - \bar{F}(t_n))^{n-1} \int_{-\infty}^{\varepsilon s} \bar{F}(n^{1/2}(s - x))\mu(t_n)^{* (n-1)}(dx). \end{aligned}$$

Similarly to the proof of Proposition 12, we have our assertion. \square

Now let us prove Lemma 3. Since

$$\begin{aligned}
& H(n, v_n^{-1/2}s) - n \int_{-\infty}^{\varepsilon v_n^{-1/2}s} \bar{F}(n^{1/2}(s - v_n^{1/2}x))\Phi_1(x)dx \\
= & \Phi_0(v_n^{-1/2}s) - n \int_{\varepsilon v_n^{-1/2}s}^{v_n^{-1/2}s} \bar{F}(n^{1/2}(s - v_n^{1/2}x))\Phi_1(x)dx \\
& + v_n^{-1/2}n^{1/2}\Phi_1(v_n^{-1/2}s) \int_0^\infty x\mu(dx) + v_n^{-1} \frac{\Phi_2(v_n^{-1/2}s)}{2} \int_0^{n^{1/2}} x^2\mu(dx) \\
= & \Phi_0(v_n^{-1/2}s) + v_n^{-1} \frac{\Phi_2(v_n^{-1/2}s)}{2} \int_0^{n^{1/2}} x^2\mu(dx) \\
& + v_n^{-1/2}n^{1/2}\eta_1((1-\varepsilon)n^{1/2}s)\Phi_1(v_n^{-1/2}s) \\
& - v_n^{-1/2}n^{1/2} \left(\int_0^{(1-\varepsilon)n^{1/2}s} \bar{F}(z)(\Phi_1(v_n^{-1/2}s - n^{-1/2}v_n^{-1/2}z) - \Phi_1(v_n^{-1/2}s))dz \right),
\end{aligned}$$

it is easy to see that there is a $C_1 > 0$ such that for $s \geq 1$

$$|H(n, v_n^{-1/2}s) - n \int_{-\infty}^{\varepsilon v_n^{-1/2}s} \bar{F}(n^{1/2}(s - v_n^{1/2}x))\Phi_1(x)dx| \leq C_1 s \Phi_1(\varepsilon v_n^{-1/2}s).$$

Combining Proposition 19 (2) and 20, we see that there is a $C_2 > 0$ such that

$$\begin{aligned}
& |P(\sum_{k=1}^n X_k > sn^{1/2}) - n \int_{-\infty}^{\varepsilon v_n^{-1/2}s} \bar{F}(n^{1/2}(s - v_n^{1/2}x))\Phi_1(x)dx| \\
\leq & 2n(n-1)\bar{F}(s^\gamma n^{1/2})^2 + \exp(\frac{6}{s^{2\gamma}} - \frac{s}{s^\gamma}) + n\bar{F}(s^\gamma n^{1/2}) \exp(\frac{6}{s^{2\gamma}} - \frac{\varepsilon s}{2s^\gamma}) \\
& + C_2 \bar{F}((1-\varepsilon)n^{1/2}s)L(n^{1/2})^{1-\delta}
\end{aligned}$$

Hence we see that there is a $C > 0$ such that

$$\sup_{s > L(n^{1/2})^{-\beta}} (n\bar{F}(n^{1/2}s))^{-1} |P(\sum_{k=1}^n X_k > sn^{1/2}) - H(n, v_n^{-1/2}s)| \leq CL(n^{1/2})^{1-\delta}$$

Therefore by Lemma 2, we have our assertion.

Now let us prove Theorem 2.

By Theorem 4, we see that there is a $C_1 > 0$ such that

$$|P(\sum_{k=1}^n X_k > sn^{1/2}) - H(n, v_n^{-1/2}s)| \leq C_1 L(n^{1/2})^{2-\delta/2}, \quad s \geq 1.$$

Note that for any $\varepsilon > 0$, there is a $C_2 > 0$ such that $n\bar{F}(n^{1/2}s) \geq C_2^{-1}s^{-3}L(n^{1/2}) \geq C_2^{-1}L(n^{1/2})^{1+\delta/2}$ for $s \leq L(n^{1/2})^{-\delta/6}$. Hence by Lemma 2, we see that there is a $C_3 > 0$ such that

$$H(n, v_n^{-1/2}s)^{-1} \leq C_3(n\bar{F}(n^{1/2}s))^{-1} \leq C_2 C_3 L(n^{1/2})^{1+\delta/2}, \quad s \leq L(n^{1/2})^{-\delta/6}.$$

So we have

$$\sup_{s \leq L(n^{1/2})^{-\delta/6}} \left| \frac{P(\sum_{k=1}^n X_k > sn^{1/2})}{H(n, v_n^{-1/2}s)} - 1 \right| \leq C_1 C_2 C_3 L(n^{1/2})^{1-\delta}$$

From this and Lemma 3, we have Equation (2).

Equation (3) is an easy consequence of Equation (2) and Lemma 2.

8 Proof of Theorem 3

First let us assume $\limsup_{n \rightarrow \infty} (1 - v_n) \log \frac{1}{L(n^{1/2})} = 0$.

we see that

$$\begin{aligned} \Phi_0(s) - \Phi_0(v_n^{-1/2}s) &= \int_s^{v_n^{-1/2}s} \Phi_1(z) dz = \int_{s^2}^{v_n^{-1}s^2} \frac{1}{\sqrt{2\pi}} e^{-y/2} \frac{dy}{2\sqrt{y}} \\ &\leq \frac{s}{2v_1} (1 - v_n) \Phi_1(s) \\ &\leq C_0 \frac{s^2}{2v_1} (1 - v_n) \Phi_0(s) \end{aligned}$$

Let $z_n = \frac{1}{L(n^{1/2})}$, then we have $\limsup_{n \rightarrow \infty} (1 - v_n) \log z_n = 0$. Hence we have

$$\sup_{s \in [1, \sqrt{3 \log z_n}]} \left| \frac{\Phi_0(v_n^{-1/2}s) + n\bar{F}(n^{1/2}s)}{\Phi_0(s) + n\bar{F}(n^{1/2}s)} - 1 \right| \leq \frac{3C_0}{2v_1} (1 - v_n) \log z_n \rightarrow 0, \quad n \rightarrow \infty.$$

We also see that for $s > \sqrt{3 \log z_n}$,

$$\begin{aligned} &\left| \frac{\Phi_0(v_n^{-1/2}s) + n\bar{F}(n^{1/2}s)}{\Phi_0(s) + n\bar{F}(n^{1/2}s)} - 1 \right| \leq \frac{C_0}{2v_1} \frac{(1 - v_n)s^2 \Phi_0(s)}{\Phi_0(s) + n\bar{F}(n^{1/2}s)} \\ &\leq \frac{C_0}{2v_1} \frac{(1 - v_n)s^4 \Phi_0(s)}{L(n^{1/2}s)} \leq \frac{C_0^2}{2\sqrt{2\pi}v_1} s^5 \exp(-s^2/2) \frac{L(n^{1/2})}{L(n^{1/2}s)} z_n \\ &\leq \frac{C_0^2}{2\sqrt{2\pi}v_1} s^6 \exp(-s^2/2) z_n \\ &\leq \frac{C_0^2}{2\sqrt{2\pi}v_1} \sup_{s \geq \sqrt{3 \log z_n}} s^6 \exp(-s^2/6) \rightarrow 0, \quad n \rightarrow \infty. \end{aligned}$$

Hence we have $\sup_{s \in [1, \infty)} \left| \frac{\Phi_0(v_n^{-1/2}s) + n\bar{F}(n^{1/2}s)}{\Phi_0(s) + n\bar{F}(n^{1/2}s)} - 1 \right| \rightarrow 0, n \rightarrow \infty$.

Next, we assume $\limsup_{n \rightarrow \infty} (1 - v_n) \log \frac{1}{L(n^{1/2})} > 0$.

Let $y_n = (1 - v_n) \log z_n$ and $s_n = \sqrt{\log z_n}$.

Then $\limsup_{n \rightarrow \infty} y_n > 0$. Hence we see that

$$\begin{aligned} \liminf_{n \rightarrow \infty} \Phi_0(s_n)^{-1} \Phi_0(v_n^{-1/2} s_n) &= \liminf_{n \rightarrow \infty} v_n^{1/2} \Phi_1(s_n)^{-1} \Phi_1(v_n^{-1/2} s_n) \\ &\leq \liminf_{n \rightarrow \infty} \exp(-v_n^{-1} (1 - v_n) s_n^2) = \exp(-\limsup_{n \rightarrow \infty} y_n) < 1 \end{aligned}$$

and

$$\begin{aligned} \Phi_0(s_n)^{-1} n \bar{F}(n^{1/2} s_n) &\leq C_0 s_n \Phi_1(s_n)^{-1} s_n^{-2} L(n^{1/2} s_n) \\ &\leq \sqrt{2\pi} C_0 M(1) L(n^{1/2})^{1/2} \rightarrow 0, \quad k \rightarrow \infty. \end{aligned}$$

Hence we have

$$\liminf_{n \rightarrow \infty} \frac{\Phi_0(v_n^{-1/2} s_n) + n \bar{F}(n^{1/2} s_n)}{\Phi_0(s_n) + n \bar{F}(n^{1/2} s_n)} < 1.$$

Therefore we have Theorem 3.

Example

We give an example in the rest of this section.

Let $x_0 \geq 1$ and $L : [x_0, \infty) \rightarrow (0, \infty)$ be a C^2 slowly varying function satisfying

$$\int_{x_0}^{\infty} \frac{L(x)}{x} dx < \infty, \quad L(x) \rightarrow 0, x \rightarrow \infty, \quad \sup_{x \geq x_0} (|L'(x)| + |L''(x)|) < \infty.$$

Then we can find $F : \mathbb{R} \rightarrow [0, 1]$ non-decreasing C^2 function with $F(-\infty) = 0$, $F(\infty) = 1$, $\int_{\mathbb{R}} |F''(x)| dx < \infty$ and $F(x) = x^{-2} L(x)$ for sufficient large $x > 0$.

Let μ be a probability measure whose distribution function is F . Then we see that μ satisfies (A3).

Let $L(x) = (\log x)^{-1} (\log \log x)^{-1-b}$, $b > 0$ for sufficiently large $x > 0$. We can easily see that $L(x)$ satisfies the above condition. For sufficiently large $n \geq 1$, we see that

$$\begin{aligned} 1 - v_n &= \int_{n^{1/2}}^{\infty} x^2 \mu(dx) = L(n^{1/2}) + 2 \int_{n^{1/2}}^{\infty} \frac{L(x)}{x} dx = L(n^{1/2}) + \frac{2}{b} (\log \log n - \log 2)^{-b} \\ &\sim \frac{2}{b} (\log \log n)^{-b} \end{aligned}$$

Hence we have the following

Proposition 21. *Let $L(x) = (\log x)^{-1} (\log \log x)^{-1-b}$, $b > 0$ for sufficiently large $x > 0$. Then we have*

$$\limsup_{n \rightarrow \infty} (1 - v_n) \log \frac{1}{L(n^{1/2})} > 0, \quad \text{for } b \in (0, 1]$$

and

$$\lim_{n \rightarrow \infty} (1 - v_n) \log \frac{1}{L(n^{1/2})} = 0, \quad \text{for } b \in (1, \infty).$$

Therefore Equation (1) does not hold for $b \in (0, 1]$.

Reference

- [1] Borovlov, A. and K. Borovkov, Asymptotic Analysis of Random Walks : Heavy Tailed Distributions, Cambridge University Press 2008, Cambridge.
- [2] Fushiya, H., and S. Kusuoka, Uniform Estimate for Distributions of the Sum of i.i.d Random Variables with Fat Tail , J. Math. Sci. Univ. Tokyo 17 (2010), 79-121.
- [3] Nagaev, S. V., Large deviations of sums of independent random variables, Ann. Probab. 7 (1979), 745-789.
- [4] Rozovskii, L. V., Probabilities of large deviations on the whole axis, Theory Probab. Appl. 38 (1994), 53-79.

UTMS

- 2010–5 Shigeo Kusuoka: *A remark on Malliavin Calculus : Uniform Estimates and Localization.*
- 2010–6 Issei Oikawa: *Hybridized discontinuous Galerkin method with lifting operator.*
- 2010–7 Hitoshi Kitada: *Scattering theory for the fractional power of negative Laplacian.*
- 2010–8 Keiju- Sono: *The matrix coefficients with minimal K -types of the spherical and non-spherical principal series representations of $SL(3, \mathbb{R})$.*
- 2010–9 Taro Asume: *On Fatou-Julia decompositions.*
- 2010–10 Yusaku Tiba: *The second main theorem of hypersurfaces in the projective space.*
- 2010–11 Hajime Fujita , Mikio Furuta and Takahiro Yoshida: *Torus fibrations and localization of index III – equivariant version and its applications.*
- 2010–12 Nariya Kawazumi and Yusuke Kuno: *The logarithms of Dehn twists.*
- 2010–13 Hitoshi Kitada: *A remark on simple scattering theory.*
- 2010–14 Kenichi Sakamoto and Masahiro Yamamoto : Corresponding AUTHOR: *Initial value/boundary value problems for fractional diffusion-wave equations and applications to some inverse problems.*
- 2010–15 Nobuhiro Nakamura: *$Pin^-(2)$ -monopole equations and intersection forms with local coefficients of 4-manifolds.*
- 2010–16 Kenji Nakahara: *Uniform estimate for distributions of the sum of i.i.d random variables with fat tail: threshold case.*

The Graduate School of Mathematical Sciences was established in the University of Tokyo in April, 1992. Formerly there were two departments of mathematics in the University of Tokyo: one in the Faculty of Science and the other in the College of Arts and Sciences. All faculty members of these two departments have moved to the new graduate school, as well as several members of the Department of Pure and Applied Sciences in the College of Arts and Sciences. In January, 1993, the preprint series of the former two departments of mathematics were unified as the Preprint Series of the Graduate School of Mathematical Sciences, The University of Tokyo. For the information about the preprint series, please write to the preprint series office.

ADDRESS:

Graduate School of Mathematical Sciences, The University of Tokyo
3–8–1 Komaba Meguro-ku, Tokyo 153-8914, JAPAN
TEL +81-3-5465-7001 FAX +81-3-5465-7012